

Mathematics

Advanced GCE A2 7890 - 2

Advanced Subsidiary GCE AS 3890 - 2

Mark Schemes for the Units

June 2006

3890-2/7890-2/MS/R/06

OCR (Oxford, Cambridge and RSA Examinations) is a unitary awarding body, established by the University of Cambridge Local Examinations Syndicate and the RSA Examinations Board in January 1998. OCR provides a full range of GCSE, A level, GNVQ, Key Skills and other qualifications for schools and colleges in the United Kingdom, including those previously provided by MEG and OCEAC. It is also responsible for developing new syllabuses to meet national requirements and the needs of students and teachers.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by Examiners. It does not indicate the details of the discussions which took place at an Examiners' meeting before marking commenced.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the Report on the Examination.

OCR will not enter into any discussion or correspondence in connection with this mark scheme.

© OCR 2006

Any enquiries about publications should be addressed to:

OCR Publications
PO Box 5050
Annersley
NOTTINGHAM
NG15 0DL

Telephone: 0870 870 6622
Facsimile: 0870 870 6621
E-mail: publications@ocr.org.uk

CONTENTS

Advanced GCE Mathematics (7890)
Advanced GCE Pure Mathematics (7891)
Advanced GCE Further Mathematics (7892)


Advanced Subsidiary GCE Mathematics (3890)
Advanced Subsidiary GCE Pure Mathematics (3891)
Advanced Subsidiary GCE Further Mathematics (3892)

MARK SCHEME ON THE UNITS

Unit	Content	Page
4721	Core Mathematics 1	1
4722	Core Mathematics 2	7
4723	Core Mathematics 3	13
4724	Core Mathematics 4	19
4725	Further Pure Mathematics 1	25
4726	Further Pure Mathematics 2	31
4727	Further Pure Mathematics 3	35
4728	Mechanics 1	41
4729	Mechanics 2	47
4730	Mechanics 3	51
4731	Mechanics 4	57
4732	Probability & Statistics 1	65
4733	Probability & Statistics 2	71
4734	Probability & Statistics 3	75
4735	Probability & Statistics 4	79
4736	Decision Mathematics 1	83
4737	Decision Mathematics 2	89
*	Grade Thresholds	97

Mark Scheme 4721
June 2006

1	(i)	$\frac{21-3}{4-1} = \frac{18}{3} = 6$	M1		Uses $\frac{y_2 - y_1}{x_2 - x_1}$
			A1	2	6 (not left as $\frac{18}{3}$)
	(ii)	$\frac{dy}{dx} = 2x + 1$	B1		
		$2 \times 3 + 1 = 7$	B1	2	
2	(i)	$27^{-\frac{2}{3}} = \frac{1}{27^{\frac{2}{3}}} = \frac{1}{9}$	M1		$\frac{1}{27^{\frac{2}{3}}}$ or $27^{\frac{2}{3}} = 9$ or 3^{-2} soi
			A1	2	$\frac{1}{9}$
	(ii)	$5\sqrt{5} = 5^{\frac{3}{2}}$	B1	1	
	(iii)	$\frac{1-\sqrt{5}}{3+\sqrt{5}} = \frac{(1-\sqrt{5})(3-\sqrt{5})}{(3+\sqrt{5})(3-\sqrt{5})}$	M1		Multiply numerator and denominator by conjugate
		$= \frac{8-4\sqrt{5}}{4}$	B1		$(\sqrt{5})^2 = 5$ soi
		$= 2 - \sqrt{5}$	A1	3	$2 - \sqrt{5}$
3	(i)	$2x^2 + 12x + 13 = 2(x^2 + 6x) + 13$	B1		$a = 2$
		$= 2[(x+3)^2 - 9] + 13$	B1		$b = 3$
		$= 2(x+3)^2 - 5$	M1		$13 - 2b^2$ or $13 - b^2$ or $\frac{13}{2} - b^2$ (their b)
			A1	4	$c = -5$
	(ii)	$2(x+3)^2 - 5 = 0$	M1		Uses correct quadratic formula or completing square method
		$(x+3)^2 = \frac{5}{2}$	A1		$x = \frac{-12 \pm \sqrt{40}}{4}$ or $(x+3)^2 = \frac{5}{2}$
		$x = -3 \pm \sqrt{\frac{5}{2}}$	A1	3	$x = -3 \pm \sqrt{\frac{5}{2}}$ or $-3 \pm \frac{1}{2}\sqrt{10}$

4	(i)	$(x-4)(x-3)(x+1)$ $\equiv (x^2 - 7x + 12)(x+1)$ $\equiv x^3 + x^2 - 7x^2 - 7x + 12x + 12$ $\equiv x^3 - 6x^2 + 5x + 12$	B1		$x^2 - 7x + 12$ or $x^2 - 2x - 3$ or $x^2 - 3x - 4$ seen
	(ii) (iii)		M1		Attempt to multiply a quadratic by a linear factor or attempt to list an 8 term expansion of all 3 brackets
			A1	3	$x^3 - 6x^2 + 5x + 12$ (AG) obtained (no wrong working seen)
			B1		+ve cubic with 3 roots (not 3 line segments)
			B1		(0, 12) labelled or indicated on y-axis
			B1	3	(-1, 0), (3, 0), (4, 0) labelled or indicated on x-axis
			M1		Reflect <i>their</i> (ii) in either x- or y-axis
			A1✓	2	Reflect <i>their</i> (ii) in x-axis
5	(i)	$1 < 4x - 9 < 5$ $10 < 4x < 14$ $2.5 < x < 3.5$	M1		2 equations or inequalities both dealing with all 3 terms
	(ii)	$y^2 \geq 4y + 5$ $y^2 - 4y - 5 \geq 0$ $(y-5)(y+1) \geq 0$ $y \leq -1, y \geq 5$	A1		2.5 and 3.5 seen oe
			A1	3	$2.5 < x < 3.5$ (or ' $x > 2.5$ <u>and</u> $x < 3.5$ ') $y^2 - 4y - 5 = 0$ soi Correct method to solve quadratic
			B1		-1, 5
			M1		(SR If both values obtained from trial and improvement, award B3)
			A1		Correct method to solve inequality
			A1	5	$y \leq -1, y \geq 5$

6	(i)	$x^4 - 10x^2 + 25 = 0$ Let $y = x^2$ $y^2 - 10y + 25 = 0$ $(y-5)^2 = 0$ $y = 5$ $x^2 = 5$ $x = \pm\sqrt{5}$	*M1 dep*M1 A1 A1	4 2	Use a substitution to obtain a quadratic or $(x^2 - 5)(x^2 - 5) = 0$ Correct method to solve a quadratic 5 (not $x = 5$ with no subsequent working) $x = \pm\sqrt{5}$ $2x^4$ or $-20x^2$ oe seen $2x^4 - 20x^2 + 50$ (integers required)
	(ii)	$y = \frac{2x^5}{5} - \frac{20x^3}{3} + 50x + 3$ $\frac{dy}{dx} = 2x^4 - 20x^2 + 50$	B1 B1	2	
	(iii)	$2x^4 - 20x^2 + 50 = 0$ $x^4 - 10x^2 + 25 = 0$ which has 2 roots	M1 A1	2	their $\frac{dy}{dx} = 0$ seen (or implied by correct answer) 2 stationary points www in any part
7	(i)	$y = x^2 - 5x + 4$ $y = x - 1$ $x^2 - 5x + 4 = x - 1$ $x^2 - 6x + 5 = 0$ $(x-1)(x-5) = 0$ $x = 1 \quad x = 5$ $y = 0 \quad y = 4$	M1 M1 A1 A1	4	Substitute to find an equation in x (or y) Correct method to solve quadratic $x = 1, 5$ $y = 0, 4$ (N.B. This final A1 may be awarded in part (ii) if y coordinates only seen in part (ii)) SR one correct (x,y) pair www B1
	(ii)	2 points of intersection	B1	1	
	(iii)	EITHER $x^2 - 5x + 4 = x + c$ has 1 solution $x^2 - 6x + (4 - c) = 0$ $b^2 - 4ac = 0$ $36 - 4(4 - c) = 0$ $c = -5$ OR $\frac{dy}{dx} = 1 = 2x - 5$ $x = 3 \quad y = -2$ $-2 = 3 + c$ $c = -5$	M1 M1 A1 A1 M1 A1 A1	4 4	$x^2 - 5x + 4 = x + c$ has 1 soln seen or implied Discriminant = 0 or $(x - a)^2 = 0$ soi $36 - 4(4 - c) = 0$ or $9 = 4 - c$ $c = -5$ Algebraic expression for gradient of curve = non-zero gradient of line used $2x - 5 = 1$ $x = 3$ $c = -5$ SR $c = -5$ without any working B1

8	(i)	Height of box = $\frac{8}{x^2}$	*B1		Area of 1 vertical face = $\frac{8}{x^2} \times x$
		4 vertical faces = $4 \times \frac{8}{x}$ $= \frac{32}{x}$	*B1		$= \frac{8}{x}$
		Total surface area = $x^2 + x^2 + \frac{32}{x}$	B1 dep on both **	3	Correct final expression
		$A = 2x^2 + \frac{32}{x}$			
	(ii)	$\frac{dA}{dx} = 4x - \frac{32}{x^2}$	B1 B1 B1	3	$4x$ kx^{-2} $-32x^{-2}$
	(iii)	$4x - \frac{32}{x^2} = 0$ $4x^3 = 32$ $x = 2$	M1 A1 M1 A1	 4	$\frac{dA}{dx} = 0$ soi $x = 2$ Check for minimum Correctly justified
					SR If $x = 2$ stated www but with no evidence of differentiated expression(s) having been used in part (iii) B1


9	(i)	$\left(\frac{4+10}{2}, \frac{-2+6}{2}\right)$ (7, 2)	M1		Uses $\left(\frac{x_1+x_2}{2}, \frac{y_1+y_2}{2}\right)$
			A1	2	(7, 2) (integers required)
	(ii)	$\sqrt{(7-4)^2 + (2-(-2))^2}$ $= \sqrt{3^2 + 4^2}$ $= 5$	M1		Uses $\sqrt{(x_2-x_1)^2 + (y_2-y_1)^2}$
			A1	2	5
	(iii)	$(x-7)^2 + (y-2)^2 = 25$	B1✓		$(x-7)^2$ and $(y-2)^2$ used (their centre)
			B1✓		$r^2 = 25$ used (their r^2)
			B1	3	$(x-7)^2 + (y-2)^2 = 25$ cao
					Expanded form: -14x and -4y used B1✓ $r = \sqrt{g^2 + f^2 - c}$ used B1✓ $x^2 + y^2 - 14x - 4y + 28 = 0$ B1 cao
	(iv)	Gradient of AB = $\frac{6-(-2)}{10-4} = \frac{4}{3}$	B1		oe
		Gradient of tangent = $-\frac{3}{4}$	B1✓		
		$y - (-2) = -\frac{3}{4}(x - 4)$	M1		Correct equation of straight line through A, any non-zero gradient
		$3x + 4y = 4$	A1	5	a, b, c need not be integers

Mark Scheme 4722
June 2006

1		$(3x-2)^4 = 81x^4 - 216x^3 + 216x^2 - 96x + 16$	M1		Attempt binomial expansion, including attempt at coeffs.
			A1 A1 A1	4 4	Obtain one correct, simplified, term Obtain a further two, simplified, terms Obtain a completely correct expansion
2	(i)	$u_2 = -1, u_3 = 2, u_4 = -1$	B1 B1	2	For correct value -1 for u_2 For correct values for both u_3 and u_4
	(ii)	Sum is $(2+(-1)) + (2+(-1)) + \dots + (2+(-1))$ i.e. $50 \times (2+(-1)) = 50$	M1 M1 A1	3 5	For correct interpretation of Σ notation For pairing, or $50 \times 2 - 50 \times 1$ For correct answer 50
	3	$y = 4x^{\frac{1}{2}} + c$ Hence $5 = 4 \times 4^{\frac{1}{2}} + c \Rightarrow c = -3$ So equation of the curve is $y = 4x^{\frac{1}{2}} - 3$	M1 A1 A1 M1 A1✓	6 6	For attempt to integrate For integral of the form $kx^{\frac{1}{2}}$ For $4x^{\frac{1}{2}}$, with or without $+c$ For relevant use of (4, 5) to evaluate c For correct value -3 (or follow through on integral of form $kx^{\frac{1}{2}}$) For correct statement of the equation in full (aef)
			A1	6	
4	(i)	Intersect where $x^2 + x - 2 = 0 \Rightarrow x = -2, 1$	M1 A1	2	For finding x at both intersections For both values correct
	(ii)	Area under curve is $\left[4x - \frac{1}{3}x^3\right]_{-2}^1$ i.e. $(4 - \frac{1}{3}) - (-8 + \frac{8}{3}) = 9$ Area of triangle is $4\frac{1}{2}$ Hence shaded area is $9 - 4\frac{1}{2} = 4\frac{1}{2}$ OR Area under curve is $\int_{-2}^1 (2 - x - x^2) dx$ $= \left[-\frac{1}{3}x^3 - \frac{1}{2}x^2 + 2x\right]_{-2}^1$ $= \left(-\frac{1}{3} - \frac{1}{2} + 2\right) - \left(\frac{8}{3} - 2 - 4\right)$ $= 4\frac{1}{2}$	M1 M1 A1 M1 A1 A1 M1 M1 A1 M1 A1 A1	6 8	For integration attempt with any one term correct For use of limits – subtraction and correct order For correct area of 9 Attempt area of triangle ($\frac{1}{2}bh$ or integration) Obtain area of triangle as $4\frac{1}{2}$ Obtain correct final area of $4\frac{1}{2}$ Attempt subtraction – either order For integration attempt with any one term correct Obtain $\pm \left[-\frac{1}{3}x^3 - \frac{1}{2}x^2 + 2x\right]$ For use of limits – subtraction and correct order Obtain $\pm 4\frac{1}{2}$ – consistent with their order of subtraction Obtain $4\frac{1}{2}$ only, following correct method only

5	(i)	$\sin^2 x = 1 - \cos^2 x \Rightarrow 2\cos^2 x + \cos x - 1 = 0$ Hence $(2\cos x - 1)(\cos x + 1) = 0$ $\cos x = \frac{1}{2} \Rightarrow x = 60^\circ$ $\cos x = -1 \Rightarrow x = 180^\circ$	M1 M1 A1 A1	4	For transforming to a quadratic in $\cos x$ For solution of a quadratic in $\cos x$ For correct answer 60° For correct answer 180° [Max 3 out of 4 if any extra answers present in range, or in radians] SR answer only is B1, B1 justification – ie graph or substitution is B2, B2
	(ii)	$\tan 2x = -1 \Rightarrow 2x = 135 \text{ or } 315$ Hence $x = 67.5^\circ \text{ or } 157.5^\circ$ OR $\sin^2 2x = \cos^2 2x$ $2\sin^2 2x = 1 \quad 2\cos^2 2x = 1$ $\sin 2x = \pm \frac{1}{2}\sqrt{2} \quad \cos 2x = \pm \frac{1}{2}\sqrt{2}$ Hence $x = 67.5^\circ \text{ or } 157.5^\circ$	M1 M1 A1 A1 M1 M1 A1 A1	4	For transforming to an equation of form $\tan 2x = k$ For correct solution method, i.e. inverse tan followed by division by 2 For correct value 67.5 For correct value 157.5 Obtain linear equation in $\cos 2x$ or $\sin 2x$ Use correct solution method For correct value 67.5 For correct value 157.5 [Max 3 out of 4 if any extra answers present in range, or in radians] SR answer only is B1, B1 justification – ie graph or substitution is B2, B2
				8	
6	(i)	(a) $100 + 239 \times 5 = \text{£}1295$	M1 A1	2	For relevant use of $a + (n-1)d$ For correct value 1295
		(b) $\frac{1}{2} \times 240 \times (100 + 1295) = \text{£}167400$	M1 A1	2	For relevant use of $\frac{1}{2}n(a+l)$ or equivalent For correct value 167400
	(ii)	$100r^{239} = 1500 \Rightarrow r = 1.01139\dots$	B1 M1 A1 M1 A1	5	For correct statement of $100r^{239} = 1500$ Attempt to find r For correct value 1.01 For relevant use of GP sum formula For correct value 124359 (3 s.f. or better)
		Hence total is $\frac{100(1.01139^{240} - 1)}{1.01139 - 1} = \text{£}124359$		9	

7	(i)	$AC^2 = 11^2 + 8^2 - 2 \times 11 \times 8 \times \cos 0.8$ $= 62.3796...$ Hence $AC = 7.90$ cm	M1 A1 A1	3	Attempt to use the cosine formula Correct unsimplified expression Show the given answer correctly
	(ii)	Area of sector $= \frac{1}{2} \times 7.90^2 \times 1.7 = 53.0$ Area of triangle $= \frac{1}{2} \times 7.90^2 \times \sin 1.7 = 30.9$ Hence shaded area $= 22.1$ cm ²	M1 M1 A1	3	Attempt area of sector using $(\frac{1}{2})r^2\theta$ Attempt area of $\triangle ACD$, using $(\frac{1}{2})r^2 \sin \theta$, or equiv Obtain 22.1
	(iii)	(arc) $DC = 7.90 \times 1.7 = 13.4$ (line) $DC^2 = 7.90^2 + 7.90^2 - 2 \times 7.90 \times 7.90 \times \cos 1.7$ $DC = 11.9$ Hence perimeter $= 25.3$ cm	M1 A1 M1 A1	4	Use $r\theta$ to attempt arc length Obtain 13.4 Attempt length of line DC using cosine rule or equiv. Obtain 25.3
				10	
8	(i)	$f(2) = 12 \Rightarrow 4a + 2b = 6$ $f(-1) = 0 \Rightarrow a - b = 12$ Hence $a = 5, b = -7$	M1 A1 M1 A1 M1 A1	6	For equating $f(2)$ to 12 For correct equation $4a + 2b = 6$ For equating $f(-1)$ to 0 For correct equation $a - b = 12$ For attempt to find a and b For both values correct
	(ii)	Quotient is $2x^2 + x - 9$ Remainder is 8	B1 M1 A1 M1 A1	5	For correct lead term of $2x^2$ For complete division attempt or equiv For completely correct quotient For attempt at remainder – either division or $f(-2)$ For correct remainder
				11	

9	(i)		M1 A1 B1	3	Attempt sketch of any exponential graph, in at least first quadrant Correct graph – must be in both quadrants For identification of (0, 1)
	(ii)	$A \approx \frac{1}{2} \times 0.5 \times \left\{ 1 + 2 \left(0.5^{\frac{1}{2}} + 0.5 + 0.5^{\frac{3}{2}} \right) + 0.5^2 \right\}$ ≈ 1.09	B1 M1 A1 A1	4	State, or imply, at least three correct y-values For correct use of trapezium rule, inc correct h For correct unsimplified expression For the correct value 1.09, or better
	(iii)	$\left(\frac{1}{2}\right)^x = \frac{1}{6} \Rightarrow x \log_{10} \frac{1}{2} = \log_{10} \frac{1}{6}$ $x = \frac{\log_{10} \frac{1}{6}}{\log_{10} \frac{1}{2}} = \frac{-\log_{10} 6}{-\log_{10} 2}$ <p>Hence $= \frac{\log_{10} 2 + \log_{10} 3}{\log_{10} 2}$</p> $= 1 + \frac{\log_{10} 3}{\log_{10} 2}$ <p>OR</p> $\left(\frac{1}{2}\right)^x = \frac{1}{6} \Rightarrow 2^x = 6$ $\Rightarrow x \log_{10} 2 = \log_{10} 6$ $x = \frac{\log_{10} 6}{\log_{10} 2}$ $= \frac{\log_{10} 2 + \log_{10} 3}{\log_{10} 2}$ $= 1 + \frac{\log_{10} 3}{\log_{10} 2}$ <p>OR</p> $\left(\frac{1}{2}\right)^x = \frac{1}{6} \Rightarrow 2^x = 6$ $2^{x-1} = 3$ $(x-1)\log_{10} 2 = \log_{10} 3$ <p>Hence $x = 1 + \frac{\log_{10} 3}{\log_{10} 2}$</p> <p>OR</p> $x = \frac{\log_{10} 2 + \log_{10} 3}{\log_{10} 2}$ $= \frac{\log_{10} 6}{\log_{10} 2}$ $x \log_{10} 2 = \log_{10} 6$ $\log_{10} 2^x = \log_{10} 6$ $2^x = 6$ $\left(\frac{1}{2}\right)^x = \frac{1}{6}$	M1 A1 M1 A1 M1 A1 M1 A1 M1 A1 M1 A1	4	For equation $\left(\frac{1}{2}\right)^x = \frac{1}{6}$ and attempt at logs Obtain $x \log\left(\frac{1}{2}\right) = \log\left(\frac{1}{6}\right)$, or equivalent For use of $\log 6 = \log 2 + \log 3$ For showing the given answer correctly For equation $2^x = 6$ and attempt at logs Obtain $x \log 2 = \log 6$, or equivalent For use of $\log 6 = \log 2 + \log 3$ For showing the given answer correctly Attempt to rearrange equation to $2^n = 3$ Obtain $2^{x-1} = 3$ For attempt at logs For showing the given answer correctly
		$x = \frac{\log_{10} 2 + \log_{10} 3}{\log_{10} 2}$ $= \frac{\log_{10} 6}{\log_{10} 2}$ $x \log_{10} 2 = \log_{10} 6$ $\log_{10} 2^x = \log_{10} 6$ $2^x = 6$ $\left(\frac{1}{2}\right)^x = \frac{1}{6}$	M1 A1 M1 A1	11	Use $\log 2 + \log 3 = \log 6$ Obtain $x \log 2 = \log 6$ Attempt to remove logarithms Show $\left(\frac{1}{2}\right)^x = \frac{1}{6}$ correctly

Mark Scheme 4723
June 2006

1	Differentiate to obtain $k(4x+1)^{-\frac{1}{2}}$ Obtain $2(4x+1)^{-\frac{1}{2}}$ Obtain $\frac{2}{3}$ for value of first derivative Attempt equation of tangent through (2, 3) Obtain $y = \frac{2}{3}x + \frac{5}{3}$ or $2x - 3y + 5 = 0$	M1 any non-zero constant k A1 or equiv, perhaps unsimplified A1 or unsimplified equiv M1 using numerical value of first derivative provided derivative is of form $k'(4x+1)^n$ A1 5 or equiv involving 3 terms
2	<u>Either:</u> Attempt to square both sides Obtain $3x^2 - 14x + 8 = 0$ Obtain correct values $\frac{2}{3}$ and 4 Attempt valid method for solving inequality Obtain $\frac{2}{3} < x < 4$ <u>Or:</u> Attempt solution of two linear equations or inequalities Obtain value $\frac{2}{3}$ Obtain value 4 Attempt valid method for solving inequality Obtain $\frac{2}{3} < x < 4$	M1 producing 3 terms on each side A1 or inequality involving $<$ or $>$ A1 M1 implied by correct answer or plausible incorrect answer A1 5 or correctly expressed equiv; allow \leq signs M1 one eqn with signs of $2x$ and x the same, second eqn with signs different A1 B1 M1 implied by correct answer or plausible incorrect answer A1 (5) or correctly expressed equiv; allow \leq signs
3	(i) Attempt evaluation of cubic expression at 2 and 3 Obtain -11 and 31 Conclude by noting change of sign (ii) Obtain correct first iterate Attempt correct process to obtain at least 3 iterates Obtain 2.34	M1 A1 A1 3 or equiv; following any calculated values provided negative then positive B1 using x_1 value such that $2 \leq x_1 \leq 3$ M1 using any starting value now A1 3 answer required to 2 d.p. exactly; 2→2.3811→2.3354→2.3410; 2.5→2.3208→2.3428→2.3401; 3→2.2572→2.3505→2.3392

4 (i) State $\ln y = (x-1)\ln 5$	B1 whether following $\ln y = \ln 5^{x-1}$ or not; brackets needed
Obtain $x = 1 + \frac{\ln y}{\ln 5}$	B1 2 AG; correct working needed; missing brackets maybe now implied
(ii) Differentiate to obtain single term of form $\frac{k}{y}$	M1 any constant k
Obtain $\frac{1}{y \ln 5}$	A1 2 or equiv involving y
(iii) Substitute for y and attempt reciprocal	M1 or equiv method for finding derivative without using part (ii)
Obtain $25 \ln 5$	A1 2 or exact equiv
5 (i) State $\sin 2\theta = 2 \sin \theta \cos \theta$	B1 1 or equiv; any letter acceptable here (and in parts (ii) and (iii))
(ii) Attempt to find exact value of $\cos \alpha$	M1 using identity attempt or right-angled triangle
Obtain $\frac{1}{4}\sqrt{15}$	A1 or exact equiv
Substitute to confirm $\frac{1}{8}\sqrt{15}$	A1 3 AG
(iii) State or imply $\sec \beta = \frac{1}{\cos \beta}$	B1
Use identity to produce equation involving $\sin \beta$	M1
Obtain $\sin \beta = 0.3$ and hence 17.5	A1 3 and no other values between 0 and 90; allow 17.4 or value rounding to 17.4 or 17.5
6 (i) <u>Either</u> : Obtain $f(-3) = -7$ Show correct process for compn of functions Obtain -47	B1 maybe implied M1 A1 3
<u>Or</u> : Show correct process for compn of functions Obtain $2 - (2 - x^2)^2$ Obtain -47	M1 using algebraic approach A1 or equiv A1 (3)
(ii) Attempt correct process for finding inverse Obtain either one of $x = \pm \sqrt{2-y}$ or both Obtain correct $-\sqrt{2-x}$	M1 as far as $x = \dots$ or equiv A1 or equiv perhaps involving x A1 3 or equiv; in terms of x now
(iii) Draw graph showing attempt at reflection in $y = x$ Draw (more or less) correct graph	M1 A1 with end-point on x -axis and no minimum point in third quadrant
Indicate coordinates 2 and $-\sqrt{2}$	A1 3 accept -1.4 in place of $-\sqrt{2}$
7 (a) Obtain integral of form $k(4x-1)^{-1}$	M1 any non-zero constant k

Obtain $-\frac{1}{2}(4x-1)^{-1}$	A1	or equiv; allow + c
Substitute limits and attempt evaluation	M1	for any expression of form $k'(4x-1)^n$
Obtain $\frac{2}{21}$	A1 4	or exact equiv
(b) Integrate to obtain $\ln x$	B1	
Substitute limits to obtain $\ln 2a - \ln a$	B1	
Subtract integral attempt from attempt at area of appropriate rectangle	M1	or equiv
Obtain $1 - (\ln 2a - \ln a)$	A1	or equiv
Show at least one relevant logarithm property	M1	at any stage of solution
Obtain $1 - \ln 2$ and hence $\ln(\frac{1}{2}e)$	A1 6	AG; full detail required
<hr/>		
8 (i) State $R = 13$	B1	or equiv
State at least one equation of form $R \cos \alpha = k$, $R \sin \alpha = k'$, $\tan \alpha = k''$	M1	or equiv; allow sin / cos muddles; implied by correct α
Obtain 67.4	A1 3	allow 67 or greater accuracy
(ii) Refer to translation and stretch	M1	in either order; allow here equiv terms such as 'move', 'shift'; with both transformations involving constants
State translation in positive x direction by 67.4	A1√	or equiv; following their α ; using correct terminology now
State stretch in y direction by factor 13	A1√ 3	or equiv; following their R ; using correct terminology now
(iii) Attempt value of $\cos^{-1}(2 \div R)$	M1	
Obtain 81.15	A1√	following their R ; accept 81
Obtain 148.5 as one solution	A1	accept 148.5 or 148.6 or value rounding to either of these
Add their α value to second value correctly attempted	M1	
Obtain 346.2	A1 5	accept 346.2 or 346.3 or value rounding to either of these; and no other solutions
<hr/>		
9 (i) Attempt to express x in terms of y	*M1	obtaining two terms

Obtain $x = e^{\frac{1}{2}y} + 1$	A1	or equiv
State or imply volume involves $\int \pi x^2$	B1	
Attempt to express x^2 in terms of y	*M1	dep *M; expanding to produce at least 3 terms
Obtain $k \int (e^y + 2e^{\frac{1}{2}y} + 1) dy$	A1	any constant k including 1; allow if dy absent
Integrate to obtain $k(e^y + 4e^{\frac{1}{2}y} + y)$	A1	
Use limits 0 and p	M1	dep *M *M; evidence of use of 0 needed
Obtain $\pi(e^p + 4e^{\frac{1}{2}p} + p - 5)$	A1 8	AG; necessary detail required
(ii) State or imply $\frac{dp}{dt} = 0.2$	B1	maybe implied by use of 0.2 in product
Obtain $\pi(e^p + 2e^{\frac{1}{2}p} + 1)$ as derivative of V	B1	
Attempt multiplication of values or expressions for $\frac{dp}{dt}$ and $\frac{dV}{dp}$	M1	
Obtain $0.2\pi(e^4 + 2e^2 + 1)$	A1✓	following their $\frac{dV}{dp}$ expression
Obtain 44	A1 5	or greater accuracy

Mark Scheme 4724
June 2006

1	$\frac{d}{dx}(xy) = x \frac{dy}{dx} + y$ $\frac{d}{dx}(y^2) = 2y \frac{dy}{dx}$	B1	s.o.i. e.g. $2x \frac{dy}{dx} + y$
	Substitute (1,2) into their differentiated equation and attempt to solve for $\frac{dy}{dx}$. [Allow subst of (2,1)]	B1	Or attempt to solve their diff equation for $\frac{dy}{dx}$ and then substitute (1,2)
	$\frac{dy}{dx} = -2$	A1	4
2	(i) $1 + (-2)(-3x) + \frac{(-2)(-3)}{1.2}(-3x)^2 (+ \dots \text{ignore})$ $= 1 + 6x$ $\dots + 27x^2$	M1	State or imply; accept $-3x^2$ & $-9x^2$
		B1	Correct first 2 terms
		A1	3 Correct third term
	(ii) $(1+2x)^2(1-3x)^{-2}$ Attempt to expand $(1+2x)^2$ & select (at least) 2 relevant products and add 55 (Accept $55x^2$) <u>SR 1</u> For expansion of $(1+2x)^2$ with 1 error, A1✓ <u>SR 2</u> For expansion of $(1+2x)^2$ & > 1 error, A0 <u>Alternative Method</u> For correct method idea of long division 1 +10x +55x ²	M1	For changing into suitable form, seen/implied
		M1	Selection may be after multiplying out
		A2✓	4 If (i) is $a+bx+cx^2$, f.t. $4(a+b)+c$
		M1	
		A1,A1,A1(4)	
3	(i) $\frac{A}{x} + \frac{B}{3-x}$ & c-u rule or $A(3-x) + Bx \equiv 3 - 2x$ $\frac{1}{x}$ $-\frac{1}{3-x}$	M1	Correct format + suitable method
		A1	seen in (i) or (ii)
		A1	3 ditto; $\frac{1}{x} - \frac{1}{3-x}$ scores 3 immediately
	(ii) $\int \frac{1}{x} (dx) = \ln x$ or $\ln x $ $\int \frac{1}{3-x} (dx) = -\ln(3-x)$ or $-\ln 3-x $ Correct method idea of substitution of limits $\ln 2 (+ \ln 1 - \ln 1) - \ln 2 = 0$ <u>Alternative Method</u> If ignoring PFs, $\ln x(3-x)$ immediately As before	B1	
		B1	Check sign carefully; do not allow $\ln(x-3)$
		M1	Dep on an attempt at integrating
		A1	4 Clearly seen; WWW AG
		B2	$\ln x(x-3) \rightarrow 0$
		M1,A1 (4)	
	(iii) Suitable statement or clear implication e.g. Equal amounts (of area) above and below (axis) or graph crosses axis or there's a root (Be lenient)	B1	1

4	(i) Working out $\mathbf{b} - \mathbf{a}$ or $\mathbf{a} - \mathbf{b}$ or $\mathbf{c} - \mathbf{a}$ or $\mathbf{a} - \mathbf{c}$ $= \pm (-3\mathbf{i} - \mathbf{j} - \mathbf{k})$ or $\pm (-2\mathbf{i} + \mathbf{j} - 2\mathbf{k})$ Method for finding magnitude of <u>any</u> vector Method for finding scalar product of <u>any</u> 2 vectors Using $\cos \theta = \frac{\mathbf{a} \cdot \mathbf{b}}{ \mathbf{a} \mathbf{b} }$ AEF for <u>any</u> 2 vectors [Alternative cosine rule method] $ \overrightarrow{BC} = \sqrt{6}$ Cosine rule used $45.3^\circ, 0.79(0), \frac{\pi}{3.97}$ (45.289378, 0.7904487)	M1 A1 M1 M1 M1 B1 M1 A1))))) ‘Recognisable’ form 6 Do not accept supplement (134.7 etc)
	(ii) Use of $\frac{1}{2} \overrightarrow{AB} \overrightarrow{AC} \sin \theta$ 3.54 (3.5355) or $\frac{5\sqrt{2}}{2}$	M1 A1	Accept $\left \frac{1}{2} \overrightarrow{AB} \times \overrightarrow{AC} \right $ 2 Accept from correct supp (134.7 etc)
5	(i) $\frac{dA}{dt}$ or kA^2 seen $\frac{dA}{dt} = kA^2$	M1 A1	 2
	(ii) Separate variables + attempt to integrate $-\frac{1}{A} = kt + c$ or $-\frac{1}{kA} = t + c$ or $-\frac{1}{A} = t + c$ Subst one of (0,0), (1,1000) or (2,2000) into eqn. Subst another of (0,),(1,1000) or (2,2000) into eqn Substitute $A = 3000$ into eqn with k and c subst $t = \frac{7}{3}$ ISW	*M1 A1 dep*M1 dep*M1 dep*M1 A1	Accept if based on $\frac{dA}{dt} = kA^2$ or A^2 Equation must contain k and/or c This equation must contain k <u>and</u> c 6 Accept 2.33, 2h 20 m
6	(i) Attempt to connect du and dx e.g. $\frac{du}{dx} = e^x$ Use of $e^{2x} = (e^x)^2$ or $(u-1)^2$ s.o.i. Simplification to $\int \frac{u-1}{u} (du)$ WWW	M1 A1 A1	But not $du = dx$ 3 AG
	(ii) Change $\frac{u-1}{u}$ to $1 - \frac{1}{u}$ or use parts $\int \frac{1}{u} du = \ln u$ Either attempt to change limits or resubstitute Show as $e+1 - \ln(e+1) - \{2 \text{ or } (1+1)\} + \ln 2$ WWW show final result as $e-1 - \ln\left(\frac{e+1}{2}\right)$	M1 A1 M1 (indep) A1 A1	If parts, may be twice if $\int \ln x dx$ is involved Seen anywhere in this part Expect new limits $e+1$ & 2 5 AG

7	(i)	Produce at least 2 of the 3 relevant eqns in λ and μ	M1	e.g. $1 + 3\lambda = -8 + \mu$, $-2 + \lambda = 2 - 2\mu$
		Solve the 2 eqns in λ & μ as far as $\lambda = \dots$ or $\mu = \dots$	M1	
		1 st solution: $\lambda = -2$ or $\mu = 3$	A1	
		2 nd solution: $\mu = 3$ or $\lambda = -2$ f.t.	A1✓	
		Substitute their λ and μ into 3 rd eqn and find 'a'	M1	
		Obtain $a = 2$ & clearly state that a cannot be 2	A1	6
	(ii)	Subst their λ or μ (& poss a) into either line eqn	M1	
		Point of intersection is $-5\mathbf{i} - 4\mathbf{j}$	A1	2 Accept any format No f.t. here
		N.B. In this question, award marks irrespective of labelling of parts		
8	(i)	<u>Integration method</u>		
		Attempt to change $\cos^2 6x$ into $f(\cos 12x)$	M1	
		$\cos^2 6x = \frac{1}{2}(1 + \cos 12x)$	A1	with $\cos^2 6x$ as the subject of the formula
		$\int = \frac{1}{2}x + \frac{1}{24}\sin 12x + c$	A1	AG Accept $\frac{1}{2}(x + \frac{1}{12}\sin 12x)$
		<u>Differentiation method</u>		
		Differentiate RHS producing $\frac{1}{2} + \frac{1}{2}\cos 12x$ ---(E)	B1	
		Attempt to change $\cos 12x$ into $f(\cos 6x)$	M1	Accept $+/- 2\cos^2 6x +/- 1$
		Simplify (E) WWW to $\cos^2 6x$ + satis finish	A1	3
<hr/>				
	(ii)	Parts with $u = x$, $dv = \cos^2 6x$	*M1	
		$x(\frac{1}{2}x + \frac{1}{24}\sin 12x) - \int(\frac{1}{2}x + \frac{1}{24}\sin 12x)dx$	A1	Correct expression only
		$\int \sin 12x dx = -\frac{1}{12}\cos 12x$	B1	Clear indication somewhere in this part
		Correct use of limits to <u>whole</u> integral	dep*M1	Accept () (-0)
		$\frac{\pi^2}{288} - \frac{\pi^2}{576} - \frac{1}{288} - \frac{1}{288}$	A1	AE unsimp exp. Accept $12x24, \sin \pi$ here
		$\frac{\pi^2}{576} - \frac{1}{144}$	+A1	6 Tolerate e.g. $\frac{2}{288}$ here
		S.R. If final marks are A0 + A0, allow SR A1 for		0.01/0.010/0.0101/0.0102/0.0101902


9	(i)	$\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}}$	M1	Used, not just quoted
		$\frac{dx}{dt} = -4 \sin t$ or $\frac{dy}{dt} = 3 \cos t$	*B1	
		$\frac{dy}{dx} = -\frac{3 \cos t}{4 \sin t}$ or $\frac{3 \cos t}{-4 \sin t}$ ISW	dep*A1	3 Also $\frac{-3 \cos t}{4 \sin t}$ provided B0 not awarded
		SR: M1 for Cartesian eqn attempt + B1 for $\frac{d}{dx}(y^2) = 2y \frac{dy}{dx}$		+ A1 as before (must be in terms of t)
<hr/>				
	(ii)	$y - 3 \sin p = \left(\text{their } \frac{dy}{dx} \right) (x - 4 \cos p)$	M1	Accept p or t here
		or $y = \left(\text{their } \frac{dy}{dx} \right) x + c$ & subst cords to find c		Ditto
		$4y \sin p - 12 \sin^2 p = -3x \cos p + 12 \cos^2 p$	A1	Correct equation cleared of fractions
		or $c = \frac{12 \sin^2 p + 12 \cos^2 p}{4 \sin p}$		
		$3x \cos p + 4y \sin p = 12$ WWW	A1	3 AG Only p here. Mixture earlier \rightarrow A0
<hr/>				
	(iii)	Subst $x = 0$ and $y = 0$ separately in tangent eqn	M1	to find R & S
		Produce $\frac{3}{\sin p}$ and $\frac{4}{\cos p}$	A1	Accept $\frac{12}{4 \sin p}$ and/or $\frac{12}{3 \cos p}$
		Use $\Delta = \frac{1}{2} \left(\frac{3}{\sin p} \cdot \frac{4}{\cos p} \right) = \frac{12}{\sin 2p}$ WWW	A1	3 AG
<hr/>				
	(iv)	Least area = 12	B1	
		$p = \frac{1}{4}\pi$ as final or only answer	B2	3 These B marks are independent.
		S.R. $45^\circ \rightarrow$ B1;		S.R. $[-12$ and e.g. $-\pi/4 \rightarrow$ B1]

4724

Mark Scheme

June 2006

Mark Scheme 4725
June 2006

1.	<p>i) $\begin{pmatrix} 7 & 4 \\ 0 & -1 \end{pmatrix}$</p> <p>(ii) $\begin{pmatrix} 3 & 0 \\ 0 & 3 \end{pmatrix}$</p> <p>$k = 3$</p>	<p>B1</p> <p>B1</p> <p>B1</p> <p>B1</p>	<p></p> <p>2</p> <p></p> <p>2</p> <p>4</p>	<p>Two elements correct</p> <p>All four elements correct</p> <p>A – B correctly found</p> <p>Find k</p>
2	<p>(i)</p>  <p>(ii) $\begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}$</p>	<p>M1</p> <p>A1</p> <p>B1 B1</p>	<p></p> <p>2</p> <p>2</p> <p>4</p>	<p>For 2 other correct vertices</p> <p>For completely correct diagram</p> <p>Each column correct</p>
3.	<p>(i) $2 + 3i$</p> <p>(ii)</p> <p>$p = -4$</p> <p>$q = 13$</p>	<p>B1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p>	<p>1</p> <p></p> <p>4</p> <p>5</p>	<p>Conjugate seen</p> <p>Attempt to sum roots or consider x terms in expansion or substitute $2 - 3i$ into equation and equate imaginary parts</p> <p>Correct answer</p> <p>Attempt at product of roots or consider last term in expansion or consider real parts</p> <p>Correct answer</p>

4.	$\Sigma r^3 + \Sigma r^2$ $\Sigma r^2 = \frac{1}{6}n(n+1)(2n+1)$ $\Sigma r^3 = \frac{1}{4}n^2(n+1)^2$ $\frac{1}{12}n(n+1)(n+2)(3n+1)$	M1 A1 A1 M1 A1		Consider the sum as two separate parts Correct formula stated Correct formula stated Attempt to factorise and simplify or expand both expressions Obtain given answer correctly or complete verification
5.	(i) $-7i$ (ii) $2+3i$ $-5+12i$ (iii) $\frac{1}{5}(4-7i)$ or equivalent	B1 B1 B1 B1 B1 M1 A1 A1	2 3 3	Real part correct Imaginary part correct iz stated or implied or $i^2 = -1$ seen Real part correct Imaginary part correct Multiply by conjugate Real part correct Imaginary part correct N.B. Working must be shown
6..	(i) Circle, Centre O radius 2 One straight line Through O with +ve slope In 1 st quadrant only (ii) $1 + \sqrt{3}$	B1 B1 B1 B1 B1 M1 A1	5 2	Sketch showing correct features Attempt to find intersections by trig, solving equations or from graph Correct answer stated as complex number

7.	<p>(i)</p> $\mathbf{A}^2 = \begin{pmatrix} 4 & 0 \\ 0 & 1 \end{pmatrix} \quad \mathbf{A}^3 = \begin{pmatrix} 8 & 0 \\ 0 & 1 \end{pmatrix}$ <p>(ii) $\mathbf{A}^n = \begin{pmatrix} 2^n & 0 \\ 0 & 1 \end{pmatrix}$</p> <p>(iii)</p>	<p>M1</p> <p>A1 A1</p> <p>B1</p> <p>B1 M1 A1 A1</p>	<p></p> <p>3</p> <p>1</p> <p>4 8</p>	<p>Attempt at matrix multiplication</p> <p>Correct \mathbf{A}^2 Correct \mathbf{A}^3</p> <p>Sensible conjecture made</p> <p>State that conjecture is true for $n = 1$ or 2 Attempt to multiply \mathbf{A}^n and \mathbf{A} or vice versa Obtain correct matrix Statement of induction conclusion</p>
8.	<p>(i)</p> $a \begin{bmatrix} a & 0 \\ 2 & 1 \end{bmatrix} - 4 \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} + 2 \begin{bmatrix} 1 & a \\ 1 & 2 \end{bmatrix}$ $a^2 - 2a$ <p>(ii)</p> $a = 0 \text{ or } a = 2$ <p>(iii) (a)</p> <p>(b)</p>	<p>M1</p> <p>A1</p> <p>A1</p> <p>M1</p> <p>A1A1ft</p> <p>B1 B1</p> <p>B1 B1</p>	<p></p> <p></p> <p>3</p> <p></p> <p>3</p> <p></p> <p>4 10</p>	<p>Correct expansion process shown</p> <p>Obtain correct unsimplified expression</p> <p>Obtain correct answer</p> <p>Solve their $\det \mathbf{M} = 0$</p> <p>Obtain correct answers</p> <p>Solution, as inverse matrix exists or \mathbf{M} non-singular or $\det \mathbf{M} \neq 0$</p> <p>Solutions, eqn. 1 is multiple of eqn 3</p>

9.	<p>(i)</p> <p>(ii)</p> <p>(iii)</p> $(n+1)^3 - 1 - \frac{3}{2}n(n+1) - n$ $\frac{1}{2}n(n+1)(2n+1)$	<p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>B1</p> <p>B1</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>A1</p>	<p>2</p> <p>2</p> <p>6</p> <p>10</p>	<p>Show that terms cancel in pairs</p> <p>Obtain given answer correctly</p> <p>Attempt to expand and simplify</p> <p>Obtain given answer correctly</p> <p>Correct Σr stated</p> <p>$\Sigma 1 = n$</p> <p>Consider sum of three separate terms on RHS</p> <p>Required sum is LHS – two terms</p> <p>Correct unsimplified expression</p> <p>Obtain given answer correctly</p>
----	---	---	---	---

10	(i) $\alpha + \beta + \gamma = 2$ $\alpha\beta\gamma = -4$	B1 B1		Write down correct values
	$\alpha\beta + \beta\gamma + \gamma\alpha = 3$	B1	3	
	(ii)	M1		Sum new roots
	$\alpha + 1 + \beta + 1 + \gamma + 1 = 5$	A1ft		Obtain numeric value using their (i)
	$p = -5$	A1ft	3	p is negative of their answer
	(iii)	M1*		Expand three brackets
		A1		$\alpha\beta\gamma + \alpha\beta + \beta\gamma + \gamma\alpha + \alpha + \beta + \gamma + 1$
		DM1		Use their (i) results
		A1ft		Obtain 2
	$q = -2$	A1ft	5	q is negative of their answer
		M2	11	Alternative for (ii) & (iii)
		A1		Substitute $x = u - 1$ in given equation
		M1		Obtain correct unsimplified equation for u
		A2		Expand
		A1 A1		Obtain $u^3 - 5u^2 + 10u - 2 = 0$
				State correct values of p and q .

Mark Scheme 4726
June 2006

- 1 Correct expansion of $\sin x$
 Multiply their expansion by $(1 + x)$
 Obtain $x + x^2 - x^3/6$
- 2 (i) Get $\sec^2 y \frac{dy}{dx} = 1$ or equivalent
 Clearly use $1 + \tan^2 y = \sec^2 y$
 Clearly arrive at A.G.
- (ii) Reasonable attempt to diff. to $\frac{-2x}{(1+x^2)^2}$
 Substitute their expressions into D.E.
 Clearly arrive at A.G.
- 3 (i) State $y = 0$ (or seen if working given)
- (ii) Write as quad. in x^2
 Use for real x , $b^2 - 4ac \geq 0$
 Produce quad. inequality in y
 Attempt to solve inequality
 Justify A.G.
- 4 (i) Correct definition of $\cosh x$ or $\cosh 2x$
 Attempt to sub. in RHS and simplify
 Clearly produce A.G.
- (ii) Write as quadratic in $\cosh x$
 Solve their quadratic accurately
 Justify one answer only
 Give $\ln(4 + \sqrt{15})$
- 5 (i) Get $(t + \frac{1}{2})^2 + \frac{3}{4}$
- (ii) Derive or quote $dx = \frac{2t}{1+t^2} dt$
 Derive or quote $\sin x = 2t/(1+t^2)$
 Attempt to replace all x and dx
 Get integral of form $A/(Bt^2+Ct+D)$
 Use complete square form as $\tan^{-1}(f(t))$
 Get A.G.
- B1 Quote or derive $x^{-1/6}x^3$
 M1 Ignore extra terms
 A1✓ On their $\sin x$; ignore extra terms; allow 3!
 SC Attempt product rule M1
 Attempt $f(0)$, $f'(0)$, $f''(0)$...
 (at least 3) M1
 Use Maclaurin accurately cao A1
- M1
 M1 May be implied
 A1
 M1 Use of chain/quotient rule
 M1 Or attempt to derive diff. equⁿ.
 A1
 SC Attempt diff. of $(1+x^2)\frac{dy}{dx} = 1$ M1, A1
 Clearly arrive at A.G. B1
- B1 Must be = ; accept x -axis; ignore any others
 M1 $(x^2y - x + (3y-1) = 0)$
 M1 Allow $>$; or $<$ for no real x
 M1 $1 \geq 12y^2 - 4y$; $12y^2 - 4y - 1 \leq 0$
 M1 Factorise/ quadratic formula
 A1 e.g. diagram / table of values of y
 SC Attempt diff. by product/quotient M1
 Solve $dy/dx = 0$ for two real x M1
 Get both $(-3, -1/6)$ and $(1, 1/2)$ A1
 Clearly prove min./max. A1
 Justify fully the inequality e.g. detailed graph B1
- B1
 M1 or LHS if used
 A1
 M1 $(2\cosh^2 x - 7\cosh x - 4 = 0)$
 A1✓ Factorise/quadratic formula
 B1 State $\cosh x \geq 1$ /graph; allow ≥ 0
 A1 cao; any one of $\pm \ln(4 \pm \sqrt{15})$ or decimal equivalent of $\ln()$
- B1 cao
 B1
 B1
 M1
 A1✓ From their expressions, $C \neq 0$
 M1 From formulae book or substitution
 A1

- 6 (i) Attempt to sum areas of rectangles
Use G.P. on $h(1+3^h+3^{2h}+\dots+3^{(n-1)h})$

Simplify to A.G.

- (ii) Attempt to find sum areas of different rect.
Use G.P. on $h(3^h+3^{2h}+\dots+3^{nh})$


Simplify to A.G.

- (iii) Get 1.8194(8), 1.8214(8) correct

- 7 (i) Attempt to solve $r=0$, $\tan \theta = -\sqrt{3}$
Get $\theta = -\frac{1}{3}\pi$ only

- (ii) $r = \sqrt{3} + 1$ when $\theta = \frac{1}{4}\pi$

- (iii)



- (iv) Formula with correct r used
Replace $\tan^2 \theta = \sec^2 \theta - 1$
Attempt to integrate their expression

Get $\theta + \sqrt{3} \ln \sec \theta + \frac{1}{2} \tan \theta$
Correct limits to $\frac{1}{4}\pi + \sqrt{3} \ln \sqrt{2} + \frac{1}{2}$

- 8 (i) Attempt to diff. using product/quotient
Attempt to solve $dy/dx = 0$
Rewrite as A.G.

- (ii) Diff. to $f'(x) = 1 \pm 2 \operatorname{sech}^2 x$
Use correct form of N-R with their expressions from correct $f(x)$
Attempt N-R with $x_1 = 2$ from previous M1
Get $x_2 = 1.9162(2)$ (3 s.f. min.)
Get $x_3 = 1.9150(1)$ (3 s.f. min.)

- (iii) Work out e_1 and e_2 (may be implied)

M1 $(h.3^h + h.3^{2h} + \dots + h.3^{(n-1)h})$

M1 All terms not required, but last term needed (or 3^{1-h}); or specify a , r and n for a G.P.

A1 Clearly use $nh = 1$

M1 Different from (i)

M1 All terms not required, but last term needed; G.P. specified as in (i), or deduced from (i)

A1

B1, B1 Allow $1.81 \leq A \leq 1.83$

M1 Allow $\pm\sqrt{3}$

A1 Allow -60°

B1, B1 AEF for r , 45° for θ

B1 Correct r at correct end-values of θ ;
Ignore extra θ used

B1 Correct shape with r not decreasing

M1 r^2 may be implied

B1

M1 Must be 3 different terms leading to any 2 of $a\theta + b \ln(\sec \theta / \cos \theta) + c \tan \theta$

A1 Condone answer $\times 2$ if $\frac{1}{2}$ seen elsewhere

A1 cao; AEF

M1

M1

A1 Clearly gain A.G.

B1 Or $\pm 2 \operatorname{sech}^2 x - 1$

M1

M1 To get an x_2

A1

A1 cao

B1 $\sqrt{-0.083(8)}$, -0.0012 (allow \pm if both of same sign); e_1 from 0.083 to 0.085

Use $e_2 \approx ke_1^2$ and $e_3 \approx ke_2^2$
 Get $e_3 \approx e_2^3/e_1^2 = -0.0000002$ (or 3)

M1
 A1 $\sqrt{\pm}$ if same sign as B1
 SC B1 only for $x_4 - x_3$

9 (i) Rewrite as quad. in e^y
 Solve to $e^y = (x \pm \sqrt{x^2 + 1})$
 Justify one solution only

M1 Any form
 A1 Allow $y = \ln(\quad)$
 B1 $x - \sqrt{x^2 + 1} < 0$ for all real x
 SC Use $C^2 - S^2 = 1$ for $C = \pm\sqrt{1+x^2}$ M1
 Use/state $\cosh y + \sinh y = e^y$ A1
 Justify one solution only B1

(ii) Attempt parts on $\sinh x$. $\sinh^{n-1}x$
 Get correct answer
 Justify $\sqrt{2}$ by $\sqrt{1+\sinh^2x}$ for $\cosh x$ when
 limits inserted
 Replace $\cosh^2 = 1 + \sinh^2$; tidy at this stage
 Produce I_{n-2}
 Gain A.G. clearly

M1
 A1 $(\cosh x \cdot \sinh^{n-1}x - \int \cosh^2 x \cdot (n-1) \sinh^{n-2}x \, dx)$
 B1 Must be clear
 M1
 A1
 A1

(iii) Attempt $4I_4 = \sqrt{2} - 3I_2$, $2I_2 = \sqrt{2} - I_0$
 Work out $I_0 = \sinh^{-1}1 = \ln(1 + \sqrt{2}) = \alpha$
 Sub. back completely for I_4
 Get $1/8(3 \ln(1+\sqrt{2}) - \sqrt{2})$

M1 Clear attempt at iteration (one at least seen)
 B1 Allow I_2
 M1
 A1 AEEF

Mark Scheme 4727
June 2006

1 (a) Identity = $1+0i$ Inverse = $\frac{1}{1+2i}$ $= \frac{1}{1+2i} \times \frac{1-2i}{1-2i} = \frac{1}{5} - \frac{2}{5}i$	B1 B1 B1 3	For correct identity. Allow 1 For $\frac{1}{1+2i}$ seen or implied For correct inverse AEFcartesian
(b) Identity = $\begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$ Inverse = $\begin{pmatrix} -3 & 0 \\ 0 & 0 \end{pmatrix}$	B1 B1 2 5	For correct identity For correct inverse
2 (a) $(z_1 z_2) = 6e^{\frac{5}{12}\pi i}$ $\left(\frac{z_1}{z_2} = \frac{2}{3}e^{-\frac{1}{12}\pi i} = \frac{2}{3}e^{\frac{23}{12}\pi i}\right)$	B1 B1 M1 A1 4	For modulus = 6 For argument = $\frac{5}{12}\pi$ For subtracting arguments For correct answer
(b) $(w^{-5}) = 2^{-5} \text{cis}\left(-\frac{5}{8}\pi\right)$ $= \frac{1}{32}\left(\cos\frac{11}{8}\pi + i\sin\frac{11}{8}\pi\right)$	M1 A1 A1 3 7	For use of de Moivre For $-\frac{5}{8}\pi$ seen or implied For correct answer (allow 2^{-5} and $\text{cis}\frac{11}{8}\pi$)

<p>3 EITHER $\mathbf{c} - \mathbf{a} = \pm[11, 3, -2]$ $(\mathbf{c} - \mathbf{a}) \times [8, 3, -6]$ $\mathbf{n} = \pm[-12, 50, 9]$ $d = \frac{ \mathbf{n} }{ [8, 3, -6] }$ $= \frac{\sqrt{2725}}{\sqrt{109}}$ $(d =) 5$</p>	<p>B1 M1* A1 ✓ M1 (dep*) A1 A1</p>	<p>For vector joining lines For attempt at vector product of $\mathbf{c} - \mathbf{a}$ and $[8, 3, -6]$ For obtaining \mathbf{n}. f.t. from incorrect $\mathbf{c} - \mathbf{a}$ For dividing \mathbf{n} by magnitude of $[8, 3, -6]$ For either magnitude correct For correct distance CAO</p>
<p>OR $\mathbf{c} - \mathbf{a} = \pm[11, 3, -2]$ $(\mathbf{c} - \mathbf{a}) \cdot [8, 3, -6]$ $\cos \theta = \pm \frac{109}{\sqrt{134}\sqrt{109}} = \pm \sqrt{\frac{109}{134}}$ $d = \sqrt{134} \sin \theta$ $(d =) 5$</p>	<p>B1 M1* A1 ✓ M1 (dep*) A1 A1</p>	<p>For vector joining lines For attempt at scalar product of $\mathbf{c} - \mathbf{a}$ and $[8, 3, -6]$ For correct $\cos \theta$ AEF. f.t. from incorrect $\mathbf{c} - \mathbf{a}$ For using trigonometry for perpendicular distance For correct expression for d in terms of θ For correct distance CAO</p>
<p>OR $\mathbf{c} - \mathbf{a} = \pm[11, 3, -2]$ $(\mathbf{c} - \mathbf{a}) \cdot [8, 3, -6]$ $x = \frac{109}{\sqrt{109}} = \sqrt{109}$ $d = \sqrt{134 - 109}$ $(d =) 5$</p>	<p>B1 M1* A1 ✓ M1 (dep*) A1 A1</p>	<p>For vector joining lines For attempt at scalar product of $\mathbf{c} - \mathbf{a}$ and $[8, 3, -6]$ For finding projection of $\mathbf{c} - \mathbf{a}$ onto $[8, 3, -6]$ f.t. from incorrect $\mathbf{c} - \mathbf{a}$ For using Pythagoras for perpendicular distance For correct expression for d For correct distance CAO</p>
<p>OR $\mathbf{CP} = \pm[-11 + 8t, -3 + 3t, 2 - 6t]$ $\mathbf{CP} \cdot [8, 3, -6] = 0$ $t = \pm 1$ OR $P = (9, 5, -1)$ $d = \sqrt{3^2 + 0^2 + 4^2}$ $(d =) 5$</p>	<p>B1 M1* A1 ✓ M1 (dep*) A1 A1 6</p>	<p>For finding a vector from $C(12, 5, 3)$ to a point on the line For using scalar product for perpendicularity For correct point. f.t. from incorrect \mathbf{CP} For finding magnitude of \mathbf{CP} For correct expression for d For correct distance CAO SR Obtain $\mathbf{CP} = [11, 3, -2] - [8, 3, -6] = \pm[3, 0, 4]$ B1 Verify $[3, 0, 4] \cdot [8, 3, -6] = 0$ M1* $d = \sqrt{3^2 + 0^2 + 4^2} = 5$ M1(dep*) A1 A1 (maximum 5 / 6)</p>


<p>4 Integrating factor $e^{\int -\frac{x^2}{1+x^3} dx}$</p> $= e^{-\frac{1}{3} \ln(1+x^3)} = (1+x^3)^{-\frac{1}{3}}$ $\Rightarrow \frac{d}{dx} \left(y(1+x^3)^{-\frac{1}{3}} \right) = \frac{x^2}{(1+x^3)^{\frac{1}{3}}}$ $\Rightarrow y(1+x^3)^{-\frac{1}{3}} = \frac{1}{2} (1+x^3)^{\frac{2}{3}} (+c)$ $\Rightarrow 1 = \frac{1}{2} + c \Rightarrow c = \frac{1}{2}$ $\Rightarrow y = \frac{1}{2} (1+x^3) + \frac{1}{2} (1+x^3)^{\frac{1}{3}}$	<p>M1</p> <p>A1</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1 ✓</p> <p>A1</p> <p>8</p>	<p>For correct process for finding integrating factor</p> <p>For correct IF, simplified (here or later)</p> <p>For multiplying through by their IF</p> <p>For integrating RHS to obtain $A(1+x^3)^k$ OR $\ln A(1+x^3)^k$</p> <p>For correct integration (+c not required here)</p> <p>For substituting (0, 1) into GS (including + c)</p> <p>For correct c. f.t. from their GS</p> <p>For correct solution. AEF in form $y = f(x)$</p>
<p>5 (i) EITHER $\mathbf{a} = [2, 3, 5]$, $\mathbf{b} = \pm[2, 2, 0]$</p> $\mathbf{n} = \mathbf{a} \times \mathbf{b} = \pm k[-10, 10, -2]$ <p>Use (2, 1, 5) OR (0, -1, 5)</p> $\Rightarrow 5x - 5y + z = 10$ <p>OR $\mathbf{a} = [2, 3, 5]$, $\mathbf{b} = \pm[2, 2, 0]$</p> <p>e.g. $\mathbf{r} = [2, 1, 5] + \lambda[2, 2, 0] + \mu[2, 3, 5]$</p> $[x, y, z] = [2 + 2\lambda + 2\mu, 1 + 2\lambda + 3\mu, 5 + 5\mu]$ $\Rightarrow 5x - 5y + z = 10$	<p>B1</p> <p>M1</p> <p>A1 ✓</p> <p>M1</p> <p>A1</p> <p>B1</p> <p>M1</p> <p>A1 ✓</p> <p>M1</p> <p>A1 5</p>	<p>For stating 2 vectors in the plane</p> <p>For finding perpendicular to plane</p> <p>For correct \mathbf{n}. f.t. from incorrect \mathbf{b}</p> <p>For substituting a point into equation $ax + by + cz = d$ where $[a, b, c] = \text{their } \mathbf{n}$</p> <p>For correct cartesian equation AEF</p> <p>For stating 2 vectors in the plane</p> <p>For stating parametric equation of plane</p> <p>For writing 3 equations in x, y, z</p> <p>f.t. from incorrect \mathbf{b}</p> <p>For eliminating λ and μ</p> <p>For correct cartesian equation AEF</p>
<p>(ii) $[2t, 3t - 4, 5t - 9]$</p>	<p>B1 1</p>	<p>For stating a point A on l_1 with parameter t AEF</p>
<p>(iii) $\pm[2t + 5, 3t - 7, 5t - 13]$</p> $\pm[2t + 5, 3t - 7, 5t - 13] \cdot [2, 3, 5] = 0$ $\Rightarrow t = 2$ $\frac{x+5}{9} = \frac{y-3}{-1} = \frac{z-4}{-3} \text{ OR}$ $\frac{x-4}{9} = \frac{y-2}{-1} = \frac{z-1}{-3}$	<p>M1</p> <p>M1</p> <p>A1</p> <p>A1 4</p> <p>10</p>	<p>For finding direction of l_2 from A and $(-5, 3, 4)$</p> <p>For using scalar product for perpendicularity with any vector involving t</p> <p>For correct value of t</p> <p>For a correct equation AEFcartesian</p> <p>SR For $2p + 3q + 5r = 0$ and no further progress award B1</p>

<p>6 (i) $(m^2 + 4 = 0 \Rightarrow) m = \pm 2i$</p> <p>CF = $A \cos 2x + B \sin 2x$</p> <p>PI = $p \sin x (+ q \cos x)$</p> <p>$-p \sin x (-q \cos x) + 4p \sin x (+4q \cos x) = \sin x$</p> <p>$\Rightarrow p = \frac{1}{3}, q = 0$</p> <p>$\Rightarrow y = A \cos 2x + B \sin 2x + \frac{1}{3} \sin x$</p>	<p>B1</p> <p>B1</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>B1 $\sqrt{6}$</p>	<p>For correct solutions of auxiliary equation (may be implied by correct CF)</p> <p>For correct CF (AEtrig but not $Ae^{2ix} + Be^{-2ix}$ only)</p> <p>State a trial PI with at least $p \sin x$</p> <p>For substituting PI into DE</p> <p>For correct p and q (which may be implied)</p> <p>For using GS = CF + PI, with 2 arbitrary constants in CF and none in PI</p>
<p>(ii) $(0, 0) \Rightarrow A = 0$</p> <p>$\frac{dy}{dx} = 2B \cos 2x + \frac{1}{3} \cos x \Rightarrow \frac{4}{3} = 2B + \frac{1}{3}$</p> <p>$A = 0, B = \frac{1}{2}$</p> <p>$\Rightarrow y = \frac{1}{2} \sin 2x + \frac{1}{3} \sin x$</p>	<p>B1 $\sqrt{}$</p> <p>M1</p> <p>A1</p> <p>A1 4</p> <p>10</p>	<p>For correct equation in A and/or B f.t. from their GS</p> <p>For differentiating their GS and substituting values for x and $\frac{dy}{dx}$</p> <p>For correct A and B Allow $A = -\frac{1}{4}i, B = \frac{1}{4}i$ from CF $Ae^{2ix} + Be^{-2ix}$</p> <p>For stating correct solution CAO</p>
<p>7 (i) $C + iS = 1 + e^{i\theta} + e^{2i\theta} + e^{3i\theta} + e^{4i\theta} + e^{5i\theta}$</p> <p>$= \frac{e^{6i\theta} - 1}{e^{i\theta} - 1}$</p> <p>$= \frac{e^{3i\theta} - e^{-3i\theta}}{e^{\frac{1}{2}i\theta} - e^{-\frac{1}{2}i\theta}} \cdot \frac{e^{3i\theta}}{e^{\frac{1}{2}i\theta}} = \frac{e^{3i\theta} - e^{-3i\theta}}{e^{\frac{1}{2}i\theta} - e^{-\frac{1}{2}i\theta}} e^{\frac{3}{2}i\theta}$</p>	<p>M1</p> <p>M1</p> <p>A1</p> <p>A1 4</p>	<p>For using de Moivre, showing at least 3 terms</p> <p>For recognising GP</p> <p>For correct GP sum</p> <p>For obtaining correct expression AG</p>
<p>(ii) $C + iS = \frac{2i \sin 3\theta}{2i \sin \frac{1}{2}\theta} \cdot e^{\frac{3}{2}i\theta}$</p> <p>Re $\Rightarrow C = \sin 3\theta \cos \frac{5}{2}\theta \operatorname{cosec} \frac{1}{2}\theta$</p> <p>Im $\Rightarrow S = \sin 3\theta \sin \frac{5}{2}\theta \operatorname{cosec} \frac{1}{2}\theta$</p>	<p>M1</p> <p>A1</p> <p>A1</p> <p>B1 4</p>	<p>For expressing numerator and denominator in terms of sines</p> <p>For $k \sin 3\theta$ and $k \sin \frac{1}{2}\theta$</p> <p>For correct expression AG</p> <p>For correct expression</p>
<p>(iii) $C = S \Rightarrow \sin 3\theta = 0, \tan \frac{5}{2}\theta = 1$</p> <p>$\theta = \frac{1}{3}\pi, \frac{2}{3}\pi$</p> <p>$\theta = \frac{1}{10}\pi, \frac{1}{2}\pi, \frac{9}{10}\pi$</p>	<p>M1</p> <p>A1</p> <p>A2 4</p> <p>12</p>	<p>For either equation deduced AEF</p> <p>Ignore values outside $0 < \theta < \pi$</p> <p>For both values correct and no extras</p> <p>For all values correct and no extras. Allow A1 for any 1 value OR all correct with extras</p>

8 (i) $r^4 \cdot a \neq a \cdot r^4$	B1 1	For stating the non-commutative product in the given table, or justifying another correct one																									
(ii) Possible subgroups order 2, 5	B1 B1 2	For either order stated For both orders stated, and no more (Ignore 1)																									
(iii) (a) $\{e, a\}$ (b) $\{e, r, r^2, r^3, r^4\}$	B1 B1 2	For correct subgroup For correct subgroup																									
(iv) order of $r^3 = 5$ $(ar)^2 = ar \cdot ar = r^4 a \cdot ar = e$ \Rightarrow order of $ar = 2$ $(ar^2)^2 = ar^2 ar \cdot r = ar^2 r^4 a \cdot r = ara \cdot r = e$ \Rightarrow order of $ar^2 = 2$	B1 M1 A1 A1 4	For correct order For attempt to find $(ar)^m = e$ OR $(ar^2)^m = e$ For correct order For correct order																									
(v) <table border="1" style="display: inline-table; vertical-align: middle;"> <tr> <th></th><th>ar</th><th>ar^2</th><th>ar^3</th><th>ar^4</th></tr> <tr> <th>ar</th><td>e</td><td>r</td><td>r^2</td><td>r^3</td></tr> <tr> <th>ar^2</th><td>r^4</td><td>e</td><td>r</td><td>r^2</td></tr> <tr> <th>ar^3</th><td>r^3</td><td>r^4</td><td>e</td><td>r</td></tr> <tr> <th>ar^4</th><td>r^2</td><td>r^3</td><td>r^4</td><td>e</td></tr> </table>		ar	ar^2	ar^3	ar^4	ar	e	r	r^2	r^3	ar^2	r^4	e	r	r^2	ar^3	r^3	r^4	e	r	ar^4	r^2	r^3	r^4	e	B1 B1 B1 B1 B1 5 14	If the border elements $ar \ ar^2 \ ar^3 \ ar^4$ are not written, it will be assumed that the products arise from that order For all 16 elements of the form e or r^m For all 4 elements in leading diagonal = e For no repeated elements in any completed row or column For any two rows or columns correct For all elements correct
	ar	ar^2	ar^3	ar^4																							
ar	e	r	r^2	r^3																							
ar^2	r^4	e	r	r^2																							
ar^3	r^3	r^4	e	r																							
ar^4	r^2	r^3	r^4	e																							

Mark Scheme 4728
June 2006

1		Momentum before = $3M - 1200 \times 3$ Momentum after = 1200×5	B1 B1	Ignore g if included; accept inconsistent directions
				(or loss of momentum of loaded wagon = $3M$ B1 gain of momentum of unloaded wagon = $1200(5 + 3)$ B1)
		$3M - 3600 = 6000$ $3(1200 + m) - 3600 = 6000$ $m = 2000$	M1 A1 A1	Equation with all terms; accept with g For any correct equation in m , M
2	(i)		M1	For resolving forces in the i direction or for relevant use of trigonometry
		$2.5 = 6.5 \sin \theta$ $\theta = 22.6^\circ$	A1 A1	AG Accept verification
	(ii)		M1	For resolving forces in the j direction or for using Pythagoras or relevant trigonometry.
		$R = 6.5 \cos 22.6^\circ$ $R = 6$	A1 A1	3

3	(i)	 <p>Time intervals 80, 40, 40 $t = 80, 120, 160$</p>	B1 B1 B1 B1 B1	<p>Line segment AB (say) of +ve slope from origin Line segment BC (say) of steeper +ve slope and shorter time interval than those for AB. SR: If the straight line segments are joined by curves, this B1 mark is not awarded Line segment CD (say) of less steep slope compared with BC.</p> <p>(An (x, t) graph is accepted and the references to more/less steep are reversed.) May be implied; any 2 correct</p>
	(ii)	Line joining $(0, 0)$ and $(160, 360)$	B1 ft	6
	(iii)	$v = 360/160$ $s = 120 + 4.5(t - 80)$ $2.25t$ $t = 106 \frac{2}{3}$ (107) SR Construction method Plotting points on graph paper t between 104 and 109 inclusive	M1 M1 A1 M1 A1 M1 A1	<p>Woman's velocity (= 2.25) For equation of man's displacement in relevant interval Accept omission of -80 Woman's displacement, awarded even if t is interpreted differently in man's expression Accept also 106.6, 106.7 but not 106</p> <p>Candidates reading the displacement intersection from graph, then dividing this distance by the woman's speed to find t, also get $v = 360/160$ M1 as above for the woman's velocity.</p>
4	(i)	Displacement is 20 m	B1	1
	(ii)	$s(t) = 0.01t^3 - 0.15t^2 + 2t$ (+A) $10 - 15 + 20 + A = 20$ Displacement is $0.01t^3 - 0.15t^2 + 2t + 5$	M1 A1 M1 A1	<p>20+c (from integration) B0</p> <p>For using $s(t) = \int v(t) dt$ Can be awarded prior to cancelling For using $s(10) = cv(20)$</p>
	(iii)	$a = 0.06t - 0.3$ $0.06t - 0.3 = 0.6$ $t = 15$ Displacement is 35 m	M1 A1 DM1 A1 B1	<p>AG</p> <p>For using $a(t) = dv/dt$ For starting solving $a(t) = 0.6$ depends on previous M1</p>

5	(i)	$R = mg$ $m = 2.55$	M1 M1 A1	3	For using $F = 5$ and $F = \mu R$ Accept 2.5 or 2.6
	(ii)a	$P \cos \alpha = 6$ $R = P \sin \alpha + 25$ $0.2R = 6$ $0.2(P \sin \alpha + 25) = 6$	B1 M1 A1ft B1 M1		For resolving vertically with 3 distinct forces Or $P \sin \alpha + (cv m)g$ For using $F = 6$ and $F = \mu R$. Can be implied by $0.2(P \sin \alpha + 25) = 6$ For an equation in $P \sin \alpha (=5)$ after elimination of R
	(ii)b	$\alpha = 39.8^\circ$ $P^2 = 6^2 + 5^2$ or $P \cos 39.8^\circ = 6$ or $P \sin 39.8^\circ = 5$	A1 M1		Accept a r t 40° For eliminating or substituting for α with cv(6). Evidence is needed that 5 is the value of $P \sin \alpha$ (rather than the original frictional force)
		$P = 7.81$	A1	8	Accept a r t 7.8
6	(i)	10500 + 3000 + 1500 Driving force below 15000 gives retardation	M1 A1	2	For summing 3 resistances Accept generalised case or specific instance
	(ii)	$35000 - 15000 = 80000a$	M1		Newton's second law for whole train
		Acceleration is 0.25 ms^{-2}	A1	2	AG Accept verification
	(iii)		M1		For applying Newton's second law to E only, at least 2 forces out of the relevant 3.
		$35000 - 10500 - 8500 = 0.25m$ Mass is 64000 kg	A1 A1	3	
	(iv)	$-15000 - 15000 = 80000a$ OR $-3000 - 10500 - 15000 = (80000 - m)a$	A1		For applying Newton's second law with all appropriate forces $a = -0.375$
		$-1500 = ma$ Mass is 4000 kg	M1 A1 A1	5	For applying Newton's second law to B only, only 1 force Or cv(a)
	(v)	$-15000 - 10500 \pm T = 64000(-0.375)$ $T = \pm 1500 \rightarrow$ forward force on E of 1500 N OR (working with A and B) $-1500 - 3000 \pm T = (80000 - 64000)(-0.375)$ $T = \pm 1500 \rightarrow$ forward force on E of 1500	B1ft B1 B1ft B1	2	Follow through cv (m_E, a), or accept use of m_E, a Follow through cv (m_E, a), or accept use of m_E, a

7	(i)	$0 = 6 + (\pm)1.5a$	M1		For using $v = u + at$ with $v = 0$
---	-----	---------------------	----	--	-------------------------------------

	$a = (\mp)4\text{ms}^{-2}$ $-mg\sin 15^\circ - F = ma$ $-0.1 \times 9.8\sin 15^\circ - F = 0.1 \times (-4)$ $R = 0.1g\cos 15^\circ$ $0.146357 \dots = \mu 0.946607$ \dots Coefficient is 0.155	A1 M1 A1 B1 M1 A1	7	For applying Newton's second law with 2 forces For using $F = \mu R$ Anything between 0.15 and 0.16 inclusive
(ii)	$mg\sin 15^\circ > \mu mg\cos 15^\circ$ (or $\tan 15^\circ > \mu$) \rightarrow particle moves down	M1 A1	2	For comparing weight component with frictional force (or \tan 'angle of friction' with μ) Awarded if conclusion is correct even though values are wrong
(iii)	$(6 + 0) \div 2 = s \div 1.5$ $s = 4.5$ $mg\sin 15^\circ - F = ma$ $0.25364 \dots - 0.146357 \dots = 0.1a$ $v^2 = 2(1.07285 \dots)4.5$ Speed is 3.11 ms^{-1}	M1 A1 M1 A1 M1 A1	6	For using $(u + v) \div 2 = s \div t$ For using Newton's second law with 2 forces Values must be correct even if not explicitly stated. Note that the correct value of friction may legitimately arise from a wrong value of μ and a wrong value of R For using $v^2 = 2as$ with any value of a Accept anything rounding to 3.1 from correct working

Mark Scheme 4729
June 2006

1		$mgh = 35 \times 9.8 \times 4$ $mgh/t = 1372/10$ 137 W	M1 A1 M1 A1	4	watch out for extras or 0.137 kW	4
2		$v^2 = 2gh$ $u = \sqrt{4g}$ or $\sqrt{39.2}$ or 6.26 $v = \sqrt{2.8g}$ or $\sqrt{27.44}$ (5.24) $I = ?$ 0.3(6.26 + 5.24) 3.45 Ns	M1 A1 A1 M1 A1✓	5	kinematics or energy speed of impact (\pm) speed of rebound (\pm) must be sum of mags. of vels. ✓ must be positive	2 5
3	(i)	$d = 2.25$ $h = 1.125$ or 1.12 or 1.13 or 9/8	B1 B1	2	3/8x6 OG (be generous) horizontal distance	7
	(ii)	$T_1 + T_2 = 12$ resolving vertically $T_1 \times 6\cos 30^\circ = 12xh$ (their h) mom(O) (their h ok for A1) $T_1 = 2.60$ N or $3\sqrt{3}/2$ $T_2 = 9.40$ N ✓ $(12 - T_1)$ above ✓ depends on at least one of the M marks ($T_s > 0$)	M1 M1 A1 A1 A1✓	5	if not then next M1 ok or mom(A) $T_2 \times 6\cos 30^\circ =$ $12(6\cos 30^\circ - h)$ or $T_2 = 9.40$ or $T_1 = 2.60$ or ✓ $(12 - T_2)$	
4	(i)	$P = 13500$ W	B1	1	or 13.5 kW	9
	(ii)	$500 = 13500/v$ $v = 27$ ms ⁻¹	M1 A1	2		
	(iii)	$15000/25 - 500 = 950a$ $a = 0.105$ or $2/19$	M1 A1 A1	3	2 parts to F A0 for 900a or 100/950	
	(iv)	$15000/26 - 500 -$ $950.9.8\sin 5^\circ = 950a$ $a = (-) .773$ ms ⁻²	M1 A1 A1	3	3 parts to F A0 for 900a s.c. accept 0.77	
5	(i)	$\bar{x} = 9$ c of m of Δ 4 cm above BD $(324 + 108)(m) \bar{y} =$ $324(m) \times 9 + 108(m) \times (18+4)$ $432 \bar{y}$ 324×9 (18 ² x 9) $108 \times (18+4)$ $\bar{y} = 12.25$	B1 B1 M1 A1 A1 A1 A1	7	ignore any working 8 cm below C/see their diagram $432 \bar{y} = 108 \times 8 + 18^2 (12 + 9)$ from C left hand side 1 st term on right hand side 2916 2 nd term on right hand side 2376 $5292 + 432$ or $49/4$	9
	(ii)	$\tan \theta = 5.75/9$ $\theta = 32.6^\circ$ or 147.4°	M1 A1✓	2	must be .../9 ✓ $\tan^{-1} ((18 - \text{their } \bar{y})/9)$ or 180° ..	

6	(i)	$T = 4.9 \text{ N}$ $T = 0.3 \times 0.2 \times \omega^2$	B1 M1 A1 A1	4	B0 for $0.5g$ or $0.3v^2/0.2$ and $\omega = v/0.2$	6
	(ii)	$\omega = 9.04 \text{ rads}^{-1}$ $\cos\theta = \sqrt{0.6/0.8} \text{ (0.968)}$ $T\cos\theta = 0.5 \times 9.8$	B1 M1 A1 A1		$(\theta=14.5^\circ)$ angle to vert. or equiv. angle consistent with diagram can be their angle	
	(iii)	$T = 5.06 \text{ N}$ $T\sin\theta = 0.5 \times v^2/0.2$ $v = 0.711 \text{ ms}^{-1}$	M1 A1 A1	3	must be a component of T ($\sin\theta = 1/4$) can be their angle	11
7	(i)	$v\sin 50^\circ$ $0 = v^2 \sin^2 50^\circ - 2 \times 9.8 \times 13$ (must be 13) $v = 20.8 \text{ ms}^{-1}$	B1 M1 A1	3	initial vertical component or $m \times 9.8 \times 13 = \frac{1}{2} m (v \sin 50^\circ)^2$	13
	(ii)	$45 = v \cos 50^\circ \cdot t$ $t = 3.36 \checkmark$ their v (3.13 for $v=22.4$) $s = v \sin 50^\circ \times t - \frac{1}{2} \times 9.8 \times t^2$ $s = -1.6$ to -2.0 inclusive (-1.68) ht above ground = 0.320 m	M1 A1 A1 A1		sin/cos mix ok for above M1 see alternative below other methods include other t_s	
	(iii)	$v_v = v \sin 50^\circ - 9.8 \times t$ $v_v = -17.0 \checkmark$ their v, t (-13.5 for 22.4) speed = $\sqrt{v_v^2 + (v \cos 50^\circ)^2}$ speed = $21.6 \text{ ms}^{-1} \checkmark$ their v and v_v (19.7 for $v = 22.4$)	M1 A1 A1 A1	4	or $v_v^2 = 2g(15 - \text{their ans to ii})$ \checkmark above for v_v or $\frac{1}{2} m v^2 - mg \times 1.68 =$ $\frac{1}{2} m \times 20.8^2$ (4 marks) M1/A1 \checkmark s, v / M1 solve/ A1 \checkmark	
	(ii)	$y = x \tan \theta - g x^2 / 2 v^2 \cos^2 \theta$ $y = 45 \tan 50^\circ -$ $9.8 \cdot 45^2 / 2 \cdot v^2 \cos^2 50^\circ$ calculate y $y = -1.6$ to -2.0 inclusive	B1 M1 A1 M1 A1		Alternative 1st 5 marks substitute v and 50° and $x=45$ can be their v should be -1.68	

8	(i)	$10 = 4 + m.x$ $e = \dots$ or rationale for $x = 2$ $m = 3$	M1 M1 A1	3	conservation of momentum	14
	(ii)	$v = 6$ $e = 4/5$ or 0.8	B1 M1 A1	3	allow sign errors for M mark watch out for lost minuses	
	(iii)	$10 - 5 = 2x + y$ ($5 = -2a + b$) $(-5 = 2c + d)$ $e = 0.8 = (y-x)/10$ $y = x + 8$ ($a + b = 8$) ($c - d = 8$) $x = -1$ ($a=1$) ($c=1$) $y = 7$ ($b=7$) ($d=-7$) $\frac{1}{2} \cdot 2.5^2 + \frac{1}{2} \cdot 1.5^2 - \frac{1}{2} \cdot 2.1^2 - \frac{1}{2} \cdot 1.7^2$ 12 J	M1 A1 M1 A1 A1 M1 A1	8	look for consistency or 1 in opp. direction to 1st K.E. lost. Must be 4 parts (37.5 – 25.5)	

± 1 in 3rd sig. fig. except where stated

Mark Scheme 4730
June 2006

1	(i)	M1		For using $I = \Delta(mv)$ in the direction of the original motion (or equivalent from use of relevant vector diagram).
	$20\cos\theta = 0.4 \times 25$ Direction at angle 120° to original motion	A1 A1	3	Accept $\theta = 60^\circ$ with θ correctly identified.
	(ii)	M1		For using $I = \Delta(mv)$ perp. to direction of the original motion (or equivalent from use of relevant vector diagram).
	$20\sin 60^\circ = 0.4v$ Speed is 43.3 ms^{-1}	A1ft A1	3	
2		M1		For applying Newton's 2 nd Law.
		M1		For using $a = v(dv/dx)$.
	$2v(dv/dx) = -(2v + 3v^2)$	A1 M1		For separating variables and attempting to integrate.
	$2/3 \ln(2 + 3v) = -x \quad (+C)$	A1ft		ft absence of minus sign,
	$[2/3 \ln 14 = C]$	M1		For using $v(0) = 4$.
	$[2/3 \ln 2 = -x + 2/3 \ln 14]$	M1		For attempting to solve $v(x) = 0$ for x .
	Comes to rest after travelling 1.30m	A1	8	AG

3	(i)		M1	For taking moments about C for the whole structure.
		$1.4R = 0.35 \times 360 + 1.05 \times 200$	A1	
		Magnitude is 240N	A1	AG
			M1	For taking moments about A for the rod AB.
		$0.7 \times 240 = 0.35 \times 200 + 1.05T$	A1	
		Tension is 93.3N	A1	6
	OR (i)		M1	For taking moments about A for AB and AC.
		$0.7R_B = 70 + 1.05T$ and $0.7R_C = 126 + 1.05T$	A1	
			M1	For eliminating T or for adding the equations, and then using $R_B + R_C = 560$.
		$0.7(560 - R_B) - 0.7R_B = 126 - 70$ or $0.7 \times 560 = 70 + 126 + 2.1T$	A1	For a correct equation in R_B only or T only
		Magnitude is 240N	A1	AG
		Tension is 93.3N	A1	6
	(ii)	Horizontal component is 93.3 N to the left	B1ft	
		$Y = 240 - 200$	M1	For resolving forces vertically.
		Vertical component is 40 N downwards	A1	3

4	(i)	M1		For using Newton's 2 nd Law perp. to string with $a = L\ddot{\theta}$.
	$L(m)\ddot{\theta} = -(m)g\sin\theta$ or $(m)\ddot{s} = -$ $(m)g\sin(s/L)$	A1		
	$\ddot{\theta} \approx -k\theta$ or $\ddot{s} = -ks$ [and motion is therefore approx. simple harmonic]	B1		
		M1		For using $T = 2\pi/\omega$ and $k =$ ω^2 or $T = 2\pi\sqrt{L/g}$ for simple pendulum.
	Period is 3.14s.	A1	5	AG
	(ii)	M1		For using $\dot{\theta}^2 = \omega^2(\theta_0^2 - \theta^2)$ or the principle of conservation of energy
	$\dot{\theta}^2 = 4(0.1^2 - 0.06^2)$ or $\frac{1}{2}m(2.45\dot{\theta})^2 =$ $2.45mg(\cos 0.06 -$ $\cos 0.1)$	A1		
	Angular speed is 0.16 rad s^{-1} .	A1	3	(0.1599... from energy method)
	OR (in the case for which (iii) is attempted before (ii))			
	(ii) $[\dot{\theta} = -0.2\sin 2t]$ $\dot{\theta} = -0.2\sin(2 \times 0.464)$ Angular speed is 0.16 rad s^{-1} .	M1 A1ft A1	3	For using $\dot{\theta} = d(A\cos nt)/dt$
	(iii)	M1		For using $\theta = A\cos nt$ or $A\sin(\pi/2 - nt)$ or for using $\theta = A\sin nt$ and $T = t_{0.1} - t_{0.06}$ ft angular displacement of 0.04 instead of 0.06
	$0.06 = 0.1\cos 2t$ or $0.1\sin(\pi/2 -$ $2t)$ or $2T = \pi/2 -$ $\sin^{-1}0.6$	A1ft		
	Time taken is 0.464s	A1	3	

5		M1	Σmv conserved in i direction.
	$2 \times 12 \cos 60^\circ - 3 \times 8 = 2a + 3b$	A1	
		M1	For using NEL
	For LHS of equation below	A1	
	$0.5(12 \cos 60^\circ + 8) = b - a$	A1	Complete equation with signs of a and b consistent with previous equation.
		M1	For eliminating a or b.
	Speed of B is 0.4 ms^{-1} in i direction	A1	
6	$a = -6.6$	A1	
	Component of A's velocity in j direction is	B1	May be shown on diagram or implied in subsequent work.
	$12 \sin 60^\circ$		
	Speed of A is 12.3 ms^{-1}	B1ft	
		M1	For using $\theta = \tan^{-1}(j \text{ comp} / \pm i \text{ comp})$
	Direction is at 122.4° to the i direction	A1ft	1 Accept $\theta = 57.6^\circ$ with
			2 θ correctly identified.
	(i) $T = 1470x/30$	B1	
	$[49x = 70 \times 9.8]$	M1	For using $T = mg$
	$x = 14$	A1	
	Distance fallen is 44m	A1ft	4
	(ii) PE loss = $70g(30 + 14)$	B1ft	
	EE gain = $1470 \times 14^2 / (2 \times 30)$	B1ft	
	$[\frac{1}{2} 70v^2 = 30184 - 4802]$	M1	For a linear equation with terms representing KE, PE and EE changes.
	Speed is 26.9 ms^{-1}	A1	4 AG
OR	(ii) $[0.5 v^2 = 14g - 68.6 + 30g]$	M1	For using Newton's 2 nd law ($vdv/dx = g - 0.7x$), integrating ($0.5 v^2 = gx - 0.35x^2 + k$), using $v(0)^2 = 60g \rightarrow k = 30g$, and substituting $x = 14$.
	For $14g + 30g$	B1ft	
	For ∓ 68.6	B1ft	
	Speed is 26.9 ms^{-1}	A1	4 Accept in unsimplified form.
			AG
	(iii) PE loss = $70g(30 + x)$	B1ft	
	EE gain = $1470x^2 / (2 \times 30)$	B1ft	
	$[x^2 - 28x - 840 = 0]$	M1	For using PE loss = KE gain to obtain a 3 term quadratic equation.
	Extension is 46.2m	A1	4
OR	(iii)	M1	For identifying SHM with $n^2 =$
			$1470 / (70 \times 30)$
		M1	For using $v_{\max} = An$
	$A = 26.9 / \sqrt{0.7}$	A1	
	Extension is 46.2m	A1	4



7	(i)	$\frac{1}{2} 0.3v^2 + \frac{1}{2} 0.4v^2$	B1		
		$\pm 0.3g(0.6\sin\theta)$	B1		
		$\pm 0.4g(0.6\theta)$	B1		
		$[0.35v^2 = 2.352\theta - 1.764\sin\theta]$	M1		For using the principle of conservation of energy.
		$v^2 = 6.72\theta - 5.04\sin\theta$	A1	5	AG
	(ii)		M1		For applying Newton's 2 nd Law radially to P and using $a = v^2/r$
		$0.3(v^2/0.6) = 0.3g\sin\theta - R$	A1		
		$[\frac{1}{2}(6.72\theta - 5.04\sin\theta) =$	M1		For substituting for v^2 .
		$0.3g\sin\theta - R]$			
		Magnitude is $(5.46\sin\theta - 3.36\theta)\text{N}$	A1		AG
	(iii)	$[5.46\cos\theta - 3.36 = 0]$	M1		For using $dR/d\theta = 0$
		Value of θ is 0.908	A1	6	
		$[T - 0.3g\cos\theta = 0.3a]$	M1		For applying Newton's 2 nd Law tangentially to P
		$[0.4g - T = 0.4a]$	M1		For applying Newton's 2 nd Law to Q
					[If $0.4g - 0.3g\cos\theta = 0.3a$ is seen, assume this derives from $T - 0.3g\cos\theta = 0.3a$ M1 and $T = 0.4g$ M0]
OR	(iii)	Component is $5.6 - 4.2\cos\theta$	A1	3	
OR	(iii)	$0.4g - 0.3g\cos\theta = (0.3 + 0.4)a$	B2		
		Component is $5.6 - 4.2\cos\theta$	B1	3	
OR	(iii)	$[2v(dv/d\theta) = 6.72 - 5.04\cos\theta]$	M1		For differentiating v^2 (from (i)) w.r.t. θ
		$2(0.6a) = 6.72 - 5.04\cos\theta$	M1		For using $v(dv/d\theta) = ar$
		Component is $5.6 - 4.2\cos\theta$	A1	3	

Mark Scheme 4731
June 2006

1	$\int x \rho \, dx = \int_0^a k(a+2x)x \, dx$ $= k \left[\frac{1}{2}ax^2 + \frac{2}{3}x^3 \right]_0^a \quad (= \frac{7}{6}ka^3)$ $\int \rho \, dx = k \int_0^a (a+2x) \, dx = k \left[ax + x^2 \right]_0^a$ $= 2ka^2$ $\bar{x} = \frac{\frac{7}{6}ka^3}{2ka^2}$ $= \frac{7}{12}a$	M1 A1 B1 M1 A1	for $\int \dots (a+2x)x \, dx$ for $\dots \left[ax + x^2 \right]_0^a$ Dependent on first M1 Accept 0.583a 5
2 (i)	$I = \frac{1}{2} \times 8 \times 0.15^2 \quad (= 0.09 \, \text{kg m}^2)$	B1	
	Using $\omega_2^2 = \omega_1^2 + 2\alpha\theta$ $25^2 = 10^2 + 2\alpha \times 75$ $\alpha = 3.5 \, \text{rad s}^{-2}$ Couple is $I\alpha = 0.09 \times 3.5$ $= 0.315 \, \text{N m}$	M1A1 M1 A1 ft	ft from wrong I and / or α , but ft requires M1M1 5
	OR Increase in KE is $\frac{1}{2} \times 0.09 \times (25^2 - 10^2)$ M1A1 ft $= 23.625 \, \text{J}$ Couple is $\frac{23.625}{75} = 0.315 \, \text{N m}$	M1 A1 ft	WD by couple is $L \times 75$ ft requires M1M1
(ii)	By conservation of angular momentum $(0.09 + I_2) \times 9 = 0.09 \times 25$ $I_2 = 0.16 \, \text{kg m}^2$	M1 A1 ft A1	Using angular momentum 3

3	$\int_1^2 \frac{1}{x^2} dx = \left[-\frac{1}{x} \right]_1^2$ $= \frac{1}{2}$ <p>Mass per unit area $\rho = 48 \text{ kg m}^{-2}$</p> $I = \int \frac{4}{3} (\rho y \delta x) \left(\frac{1}{2} y \right)^2$ $= \int \frac{1}{3} \rho y^3 dx$ $= \frac{1}{3} \rho \int_1^2 \frac{1}{x^6} dx$ $= \frac{1}{3} \rho \left[-\frac{1}{5x^5} \right]_1^2$ $= \frac{31}{480} \rho = \frac{31}{480} \times 48$ $= 3.1 \text{ kg m}^2$	M1 A1 B1 M1 A1 A1 ft A1 A1	 For integral of y^3 For correct integration of $\frac{1}{x^6}$ 8
4 (i)	$RC = 2a \cos \theta$ $EPE = \frac{5mg}{2a} (2a \cos \theta)^2$ $GPE = mga \sin 2\theta + 2mg(2a \sin 2\theta)$ $V = 10mga \cos^2 \theta + 5mga \sin 2\theta$ $\frac{dV}{d\theta} = -20mga \cos \theta \sin \theta + 10mga \cos 2\theta$ $= -10mga \sin 2\theta + 10mga \cos 2\theta$ <p>For equilibrium, $10mga(\cos 2\theta - \sin 2\theta) = 0$</p> $\tan 2\theta = 1$ $\theta = \frac{1}{8} \pi$	B1 M1 M1 A1 B1 M1 A1	or $RC^2 = 2a^2 + 2a^2 \cos 2\theta$ One term sufficient for M1 Correct differentiation of $\cos^2 \theta$ (or $\cos 2\theta$) and $\sin 2\theta$ For using $\frac{dV}{d\theta} = 0$ Accept $22\frac{1}{2}^\circ$, 0.393 7
(ii)	$\frac{d^2V}{d\theta^2} = -20mga \cos 2\theta - 20mga \sin 2\theta$ <p>When $\theta = \frac{1}{8} \pi$, $\frac{d^2V}{d\theta^2} (= -20\sqrt{2} mga) < 0$</p> <p>Hence the equilibrium is unstable</p> <hr/> <p>OR Other method for determining whether V has a maximum or a minimum</p> <p>Correct determination</p> <p>Equilibrium is unstable</p>	B1 ft M1 A1 M1 A1 ft A1	 Determining the sign of V'' Correctly shown 3 Correctly shown

5 (i)	$I = \frac{1}{3}(20)(0.3^2 + 0.9^2) + 20 \times 0.9^2$ $= 22.2 \text{ kg m}^2$	M1 M1 A1 (ag)	MI of lamina about any axis Use of parallel (or perp) axes rule
	OR $I = \frac{1}{3} \times 20 \times 0.3^2 + \frac{4}{3} \times 20 \times 0.9^2$ $= 22.2 \text{ kg m}^2$	M1M1 A1	Correctly obtained As above
(ii)	Total moment is $20 \times 9.8 \times 0.9 \cos \theta - 44.1$ Angular acceleration is zero when moment is zero $\cos \theta = \frac{44.1}{20 \times 9.8 \times 0.9} = 0.25$	M1 M1 A1 (ag)	
(iii)	Maximum angular speed when $\cos \theta = 0.25$ $\theta = 1.318$ Work done against couple is 44.1×1.318 By work energy principle, $\frac{1}{2} I \omega^2 = 20 \times 9.8 \times 0.9 \sin \theta - 44.1 \theta$ $\omega = 3.19 \text{ rad s}^{-1}$	M1 A1 M1 A1 ft A1	Equation involving work, KE and PE

<p>6 (i)</p>	<p>As viewed from P</p>  $\sin \alpha = \frac{1790}{7400}$ $\alpha = 14.0^\circ$ <p>Bearing of relative velocity is $50 - \alpha = 036^\circ$ or $50 + \alpha = 064^\circ$</p>	<p>M1</p> <p>A1 (ag) B1 ft</p> <p>3</p>	<p>For 64 or ft $50 + \alpha$</p>
<p>(ii)</p>	<p>Velocity diagram</p>  $\frac{\sin \beta}{7} = \frac{\sin 106}{10}$ $\beta = 42.3^\circ$ <p>Bearing of v_Q is $36 + \beta = 078.3^\circ$</p>	<p>B1</p> <p>M1</p> <p>A1 A1</p> <p>4</p>	<p>Correct diagram (may be implied)</p> <p>Correct triangle must be intended</p> <p>Accept 78°</p>
<p>(iii)</p>	$\frac{w}{\sin 31.7} = \frac{10}{\sin 106}$ $w = 5.47 \text{ ms}^{-1}$	<p>M1</p> <p>A1</p> <p>2</p>	<p>If cosine rule is used, M1 also requires an attempt at solving the quadratic</p>
	<p>Alternative for (ii) and (iii)</p> $\begin{pmatrix} w \sin 36 \\ w \cos 36 \end{pmatrix} = \begin{pmatrix} 10 \sin \theta \\ 10 \cos \theta \end{pmatrix} - \begin{pmatrix} 7 \sin 110 \\ 7 \cos 110 \end{pmatrix}$ <p>Obtaining an equation in θ only, and solving it M1</p> <p>$\theta = 78.3^\circ$ A2</p> <p>Obtaining an equation in w only, and solving it M1</p> <p>$w = 5.47 \text{ ms}^{-1}$ A1</p>	<p>B1</p>	<p>e.g. $10 \sin \theta - 7.2654 \cos \theta = 8.3173$ or A1A1 if another angle found first</p>

4731

Mark Scheme

June 2006

(iv)	$QC = \sqrt{7400^2 - 1790^2} = 7180 \text{ m}$ Time taken is $\frac{7180}{5.468}$ $= 1310 \text{ s}$	M1 M1 A1 ft	3 (Or M2 for other complete method for finding the time) For attempt at relative distance $\div w$ (not awarded for $7400 \div w$) or 21.9 minutes ft is $7180 \div w$
(v)	Bearing of CP is $90 + 36 = 126^\circ$	B1 1	

7 (i)	$I = \frac{1}{3}m(3a)^2 + m(2a)^2$ $= 7ma^2$ $mg(2a \sin \theta) = I\alpha$ $\alpha = \frac{2g \sin \theta}{7a}$	M1 A1 M1 A1	Using parallel axes rule
		4	
	(ii) By conservation of energy $\frac{1}{2}I\omega^2 = mg(2a \cos \frac{1}{3}\pi - 2a \cos \theta)$ $\frac{7}{2}ma^2\omega^2 = mga(1 - 2 \cos \theta)$ $\omega = \sqrt{\frac{2g(1 - 2 \cos \theta)}{7a}}$	M1 A1 A1 (ag)	Equation involving KE and PE Need to see how $\frac{1}{3}\pi$ is used Correctly obtained
		3	
(iii)	$mg \cos \theta - R = m(2a\omega^2)$ $R = mg \cos \theta - \frac{4}{7}mg(1 - 2 \cos \theta)$ $= \frac{1}{7}mg(15 \cos \theta - 4)$	M1 A1 A1	For radial acceleration $r\omega^2$
	$mg \sin \theta - S = m(2a\alpha)$ $S = mg \sin \theta - \frac{4}{7}mg \sin \theta$ $= \frac{3}{7}mg \sin \theta$	M1 A1 A1	For transverse acceleration $r\alpha$
		6	
	OR $S(2a) = I_G\alpha = (3ma^2)\alpha$ $S = \frac{3}{7}mg \sin \theta$	M1A1 A1	Must use I_G
(iv)	When $\cos \theta = \frac{1}{3}$, $\sin \theta = \frac{\sqrt{8}}{3}$, $\tan \theta = \sqrt{8}$ $R = \frac{1}{7}mg$, $S = \frac{\sqrt{8}}{7}mg$ Angle with R is $\tan^{-1} \frac{S}{R} = \tan^{-1} \sqrt{8} = \theta$ so the resultant force is vertical Magnitude is $\sqrt{R^2 + S^2}$ $= \frac{1}{7}mg\sqrt{1+8} = \frac{3}{7}mg$	M1 A1 M1 A1	
		4	
	OR When resultant force is F vertically upwards $S = F \sin \theta$, hence $F = \frac{3}{7}mg$ $R = F \cos \theta$, so $\frac{1}{7}mg(15 \cos \theta - 4) = \frac{3}{7}mg \cos \theta$ $\cos \theta = \frac{1}{3}$	M1A1 M1 A1	
	OR Horizontal force is $R \sin \theta - S \cos \theta$ $= \frac{1}{7}mg(15 \cos \theta - 4) \sin \theta - \frac{3}{7}mg \sin \theta \cos \theta$ $= \frac{1}{7}mg \sin \theta (12 \cos \theta - 4)$ $= 0$ when $\cos \theta = \frac{1}{3}$	M1 A1	

4731

Mark Scheme

June 2006

	<p>Vertical force is $R \cos \theta + S \sin \theta$ $= \frac{1}{7}mg \times \frac{1}{3} + \frac{3}{7}mg \times \frac{8}{9} = \frac{3}{7}mg$</p>	M1A1	
--	--	------	--

Mark Scheme 4732
June 2006

Note: "(3 sfs)" means "answer which rounds to ... to 3 sfs". If correct ans seen to ≥ 3 sfs, ISW for later rounding
 Penalise 2 sfs only once in paper.

1(i)	Negative, because (grad or coeff of x in 1 st eqn or x -value or reg coeff or B or -0.6) is negative	B1	1	Neg because x incr & y decr
(ii)	$x = -1.6 \times 7.0 + 21$ $x = 9.8$	M1 A1	2	Sub $y=7.0$ in 2 nd eqn. Allow 1 sign error If sub in both must choose 2nd
(iii)	$y = -0.6(-1.6y + 21) + 13$ or similar $\bar{x} = 5, \bar{y} = 10$	M1 A1A1	3	Obtain correct eqn in 1 variable. Allow 1 num'l error Allow without bars
Total			6	
In qus 2 & 3 "prod" means "product of two probabilities"				
2(i)	$^4/7$ or 0.571 (3 sfs)	B1	1	
(ii)	$^5/8 \times ^4/7 + ^3/8 \times ^5/8$ $= ^{265}/_{448}$ or 0.592 (3 sfs)	M1M1 A1	3	M1: one correct prod or add any two prods M1: all correct
(iii)	$^3/8 \times ^5/8 + ^5/8 \times ^3/7$ $= ^{225}/_{448}$ or 0.502 (3 sfs)	M1M1 A1	3	M1: one correct prod or add any two prods M1: all correct
Total			7	
3(i)	$\frac{7!}{3! \times 2(1)!}$ $= 420$	M1M1 A1	3	M1: $7!/(a \text{ factorial})$; or $\dots \div (3! \times 2(1)!)$ M1: all correct
(ii)	$\frac{5!}{2(1)!}$ $= 60$	M1 A1	2	M1: $5!$ seen (not part of a C) or $5 \times 4!$ or 120 seen or $\dots \div 2(1)!$ alone
(iii)	$1 - ^4/7 \times ^3/6$ or $1 - ^4C_2 / ^7C_2$ or $1 - ^4P_2 / ^7P_2$ or $^3/7 \times ^2/6 + ^3/7 \times ^4/6 + ^4/7 \times ^3/6$ oe or $^3C_2 / ^7C_2 + ^3C_1 \times ^4C_1 / ^7C_2$ $= ^5/7$ or 0.714 (3 sfs)	M1M1 A1	3	M1: $1 - \text{prod}$ or $1 - \dots / ^7C_2$ or $1 - ^4C_2 / ..$ (or Ps) or add 3 prods or add 2 correct prods or $^3C_2 / ^7C_2$ or $^3C_1 \times ^4C_1 / ^7C_2$ or add ≥ 5 out of 7 correct prods M1: all correct
Total			8	

4(i)	0.4207 or 0.421 (3 sfs) or $0.8^{25} + 25 \times 0.8^{24} \times 0.2 + \dots {}^{25}C_4 \times 0.4^{21} \times 0.2^4$ 0.579(3)	B1 B1 2	or $1 - 0.6167$ or 0.3833 (3 sfs) or $1 - (6 \text{ correct terms, } 0 \text{ to } 5)$
(ii)	${}^{10}C_3 \times (1-0.27)^7 \times 0.27^3$ $= 0.261$ (3 sfs)	M1 A1 2	
(iii)	<div> $0.73^9 = 0.059$ $0.73^{10} = 0.043$ $n = 10$ </div> <div> Allow "=" thro'out $1 - 0.73^n > 0.95$ or $0.73^n < 0.05$ $n \log 0.73 < \log 0.05$ oe </div>	M1 M1 A1 3	or $1 - {}^nC_0 \times 0.27^0 \times 0.73^n > 0.95$ oe allow incorrect sign M1 must be correct ft $(1 - 0.27)$ from (ii) for M1M1 10 with incorrect sign in wking: SCB2 10 with just $0.73^9 = 0.059$: M1M1A1
Total		7	
5(i)	$\frac{1}{3} + \frac{1}{4} + p + q = 1$ oe $0 \times \frac{1}{3} + 1 \times \frac{1}{4} + 2p + 3q = 1\frac{1}{4}$ oe equalize coeffs, eg mult eqn (i) by 2 or 3 Or make p or q subject of (i) or (ii) $p = \frac{1}{4}, q = \frac{1}{6}$ oe	B1 B1 M1 A1A1 5	allow one error. ft their equns subst or subtr not nec'y
(ii)	$\sum x^2 p$ (not $\frac{1}{4}$ or $\frac{1}{3}$ etc) $(= 2\frac{3}{4})$ $- (\frac{1}{4})^2$ $= 1.1875$ or $1\frac{3}{16}$ oe $sd = \sqrt{(\text{their } 1.1875)} = 1.09$ (3 sfs)	M1 M1 A1 B1f 4	≥ 2 non-zero terms correct. dep +ve result indep if +ve result or $\square x - 1\frac{1}{4})^2 p$ $(\geq 2 \text{ (non-0) terms correct): M2}$ ft (i) ($0 \leq p, q < 1$) or letters p, q both M1s cao dep 1st M1 & / (+ve no.) eg $\sqrt{2.75} = 1.66$
Total		9	

6(i)(a)	Ranks: 2 4 7 5 3 1 6 6 4 1 3 5 7 2 7 1 6 3 2 5 4 1 7 2 5 6 3 4 $\sum d^2$ (= 60) $r_s = 1 - \frac{6 \times 60}{7 \times 48}$ = $-1/14$ or -0.071 (3 dps)	M1 A1 M1 M1 A1 5	≥ 5 ranks correct in each set all correct dep ranks attempted even if opp orders, allow arith errors Correct formula with $n = 7$, dep 2 nd M1 calc r for ranks: $S_{xx}=S_{yy}= 140 - 28^2/7$ $S_{xy}= 110-28^2/7$ (= 28) (= -2) corr subst in one corr S (any version):M1 corr subst in $r = S_{xy} / \sqrt{(S_{xx}S_{yy})}$:M1 -0.07 without wking: M1A1M2A0 No mks unless $ r_s \leq 1$ fit their r_s Must refer to context. Not "little corr'n between dist and com" not "strong disagreement" Ignore other comment
(b)	Little (or no) connection (agreement, rel'nship) between dist and commission Allow disagreement	B1ft 1	
(c)	Unchanged. No change in rank	B1B1 2	
(ii)(a)	= -1	B1 1	indep
(b)	Close to -1 or, eg ≈ -0.9	B1	cao not referring to "corr'n" rather than r allow "neg", not neg corr'n or neg skew
Total		10	

7(i)	<p>Midpoints attempted ≥ 2 classes $\sum xf / 100$ or $\sum xf / \sum f$ attempted ≥ 2 terms x within class, not class width Mean = 27.2 (to 3 sfs) (not 27.25) art 27.2 from fully correct wking</p> <p>$\sum x^2 f$ or $\sum (x - \bar{x})^2 f \geq 2$ terms $\sqrt{(\sum x^2 f / 100 - \bar{x}^2)}$ or $\sqrt{((\sum x - \bar{x})^2 f / 100)}$ or $\sqrt{\sum f}$ fully corr method, not $\sqrt{\text{neg}}$ = 40.5 to 41.1 (3 sfs)</p>	<p>M1 M1 A1 M1 M1 A1</p>	<p>Correct (149.5) 2720.5/100 27.2 240702.25 40.82</p>	<p>With 150 2725/100 27.25 242050 40.96</p>	<p>Tot = 2000 Allow Ms & poss As</p>
(ii)	<p>Recog LQ in 1st class & UQ in 3rd class</p> <p><u>Graph:</u> Attempt 25(.25)th value <u>Interp:</u> LQ = 3.0 to 4.3 Attempt 75(.75)th value UQ = 27 to 29</p> <p>Subtract IQR = 23 or 24 or 25</p>	<p>B1 M1 M1 A1</p>	<p>6 both nec'y dep B1 or M1 integer. dep M2</p>		
(iii)(a)	Increase	B1	1		
(b)	Increase	B1	1		
(c)	No change	B1	1		Ignore "probably" etc
Total			13		
8(i)	Geometric. Each attempt (or result or try) indep	B1 B1	2		In context. Not "events, trials, outcomes". Ignore extra
(ii)(a)	$(\frac{2}{3})^3 \times \frac{1}{3}$ = $\frac{8}{81}$ or 0.0988 (3 sfs)	M2 A1	3		$(\frac{2}{3})^2 \times \frac{1}{3}$ or $(\frac{2}{3})^4 \times \frac{1}{3}$: allow other numerical "p" ($0 < p < 1$): M1
(b)	$(\frac{2}{3})^3$ $1 - (\frac{2}{3})^3$ = $\frac{19}{27}$ or 0.704 (3sfs)	M1 M1 A1	3		not $(\frac{2}{3})^3 \times \dots$ or $\frac{1}{3} + \frac{2}{3} \times \frac{1}{3} + (\frac{2}{3})^2 \times \frac{1}{3}$ M2 $1 - (\frac{2}{3})^4$ or $1 - ("q")^4$ M1 or 3 terms, with 2 correct M1 or 3 correct terms + 1 extra M1 or "p" + "qp" + "q ² p" M1 or 1 - sum of 3 correct terms M1 <p>"p" means num value, not $\frac{1}{3}$</p>
(iii)	3	B1f	1		or $\frac{1}{p}$
(iv)	$1 - \frac{19}{27}$ $(\frac{8}{27})^2 \times \frac{19}{27}$ = $\frac{1216}{19683}$	<p>(1 - 0.7037) or 0.2963 0.2963² x 0.7037 = 0.0618 (3 sfs)</p> <p>M1 M1 A1</p>	<p>3</p>		<p>ft (b) for M1M1 must see method if ft Allow figs rounded to 2 sfs for M1M1 cao. allow art 0.0618 or 0.0617</p>
Total			12		

Total 72 marks

Mark Scheme 4733
June 2006

1	$\mu = \frac{3}{37} \int_3^4 x^3 dx = \frac{3}{37} \left[\frac{x^4}{4} \right]_3^4$ $= 12 \frac{123}{185} \text{ or } 12.665$ $\sigma^2 = 12 \frac{123}{185} - 3 \left(\frac{81}{148} \right)^2 = \mathbf{0.0815}$	M1 M1 A1 A1 M1 A1 6	Integrate $xf(x)$, limits 3 & 4 <i>[can be implied]</i> $\left[\frac{525}{148} \text{ or } 3.547 \right]$ Attempt to integrate $x^2f(x)$, limits 3 & 4 Correct indefinite integral, any form $\frac{2343}{185}$ or in range [12.6, 12.7] <i>[can be implied]</i> Subtract their μ^2 Answer, in range [0.0575, 0.084]
2	(i) Find $P(R \geq 6)$ or $P(R < 6)$ $= 0.0083$ or 0.9917 Compare with 0.025 [can be from N] [0.05 if "empty LH tail stated"] Reject H_0 (ii) $n = 9$, $P(\leq 1) = 0.0385$ [> 0.025] $n = 10$, $P(\leq 1) = 0.0233$ [< 0.025] Therefore $n = 9$	M1 A1 B1 A1✓ 4 M1 A1 B1 3	Find $P(= 6)$ from tables/calc, OR RH critical region $P(\geq 6)$ in range [0.008, 0.0083] or $P(< 6) = 0.9917$ OR CR is 6 with probability 0.0083/0.9917 Explicitly compare with 0.025 [or 0.975 if consistent] OR state that result is in critical region Correct comparison and conclusion, ✓ on their p At least one, or $n = 8$, $P(\leq 1) = 0.0632$ Both of these probabilities seen, don't need 0.025 Answer $n = 9$ only, indep't of M1A1, <i>not</i> from $P(= 1)$
3	(i) $(140 - \mu)/\sigma = -2.326$ $(300 - \mu)/\sigma = 0.842$ Solve to obtain: $\mu = \mathbf{257.49}$ $\sigma = \mathbf{50.51}$ (ii) Higher as there is positive skew	M1 B1 A1✓ M1 A1 A1 6 B1 B1 2	One standardisation equated to Φ^{-1} , allow "1-", σ^2 Both 2.33 and 0.84 at least, ignore signs Both equations completely correct, ✓ on their z Solve two simultaneous equations to find one variable μ value, in range [257, 258] σ in range [50.4, 50.55] "Higher" or equivalent stated Plausible reason, allow from normal calculations
4	(i) Each element equally likely to be selected (and all selections independent) OR each possible sample equally likely (ii) $B(6, 5/8)$ ${}^6C_4 p^4 (1-p)^2$ $= \mathbf{0.32187}$ (iii) $N(37.5, 225/16)$ $\frac{39.5 - 37.5}{3.75} = 0.5333$ $1 - \Phi(0.5333)$ $= \mathbf{0.297}$	B1 1 M1 M1 A1✓ 3 B1 B1 M1 dep A1 dep M1 A1 6	One of these two. "Selections independent" alone is insufficient, but don't need this. An example is insufficient. $B(6, 5/8)$ stated or implied, allow e.g. 499/799 Correct formula, any p Answer, a.r.t. 0.322, can allow from wrong p Normal, mean 37.5, or 37.47 from 499/799, 499/800 14.0625 or 3.75 seen, allow 14.07/14.1 or 3.75 Standardise, wrong or no cc, np , npq , no ✓ n Correct cc, ✓ npq , signs can be reversed Tables used, answer < 0.5 , $p = 5/8$ Answer, a.r.t. 0.297 SR: $np < 5$: $Po(np)$ stated or implied, B1

5	(i)	B(303, 0.01) $\approx \text{Po}(3.03)$	B1 B1	2	B(303, 0.01) stated, allow $p = 0.99$ or 0.1 Allow Bin implied clearly by parameters Po(3.03) stated or implied, can be recovered from (ii)
	(ii)	$e^{-3.03} (1 + 3.03 + \frac{3.03^2}{2}) = 0.4165$ AG	M1 A1	2	Correct formula, ± 1 term or "1 -" or both Convincingly obtain 0.4165(02542) [Exact: 0.41535]
	(iii)	302 seats $\Rightarrow \mu = 3.02$ $e^{-3.02} (1 + 3.02 + \frac{3.02^2}{2}) = 0.1962$ $0.196 < 0.2$ So 302 seats.	M1 M1 A1 A1 A1	5	Try smaller value of μ Formula, at least one correct term Correct number of terms for their μ 0.1962 [or 0.1947 from exact] Answer 302 only
	SR: B(303, 0.99): B1B0; M0; M1 then N(298.98, 2.9898) or equiv, standardise: M1A1 total 4/9 SR: $p = 0.1$: B(303, 0.1), N(30.3, 27.27) B1B0; Standardise 2 with np & \sqrt{npq} , M1A0; N(0.1n, 0.09n); standardise with np & \sqrt{npq} ; solve quadratic for \sqrt{n} ; $n = 339$: M1M1M1A1, total SR: 6/9 B(303, 0.01) \approx N(3.03, 2.9997): B1B0; M0A0; M1A0				
6	(i)	Customers arrive independently	B1	1	Valid reason in context, allow "random"
	(ii)	$1 - 0.9921$ = 0.0079	M1 A1	2	Poisson tables, "1 -", or correct formula ± 1 term Answer, a.r.t. 0.008 [1 - 0.9384 = 0.0606: M1A0]
	(iii)	N(48, 48) $z = \frac{55.5 - 48}{\sqrt{48}}$ $= 1.0825$ $1 - \Phi(1.0825)$ = 0.1394	B1 B1 M1 dep A1 dep M1 A1	6	Normal, mean 48 Variance or SD same as mean Standardise, wrong or no cc, $\mu = \lambda$ Correct cc, $\sqrt{\lambda}$ Use tables, answer < 0.5 Answer in range [0.139, 0.14]
	(iv)	$e^{-\lambda} < 0.02$ $\lambda > -\ln 0.02$ $= 3.912$ $0.4t = 3.912$: $t = 9.78$ minutes $t = 9$ minutes 47 seconds	M1 M1 A1 M1 A1	5	Correct formula for P(0), OR P(0 $\lambda = 4$) at least ln used OR $\lambda = 3.9$ at least by T & I 3.91(2) seen OR $\lambda = 3.91$ at least by T & I Divide λ by 0.4 or multiply by 150, any distribution 587 seconds ± 1 sec [inequalities not needed]

7	(i) $\frac{c - 4000}{60 / \sqrt{50}} = 1.645$ Solve $c = 4014$ [4013.958] Critical region is > 4014	M1 B1 A1✓ M1 A1 A1✓ 6	Standardise unknown with $\sqrt{50}$ or 50 [ignore RHS] $z = 1.645$ or -1.645 seen Wholly correct eqn, $\sqrt{}$ on their z [1 – 1.645: M1B1A0] Solve to find c Value of c , a.r.t. 4014 Answer " > 4014 ", allow \geq , $\sqrt{}$ on their c , needs M1M1
	(ii) Use "Type II is: accept when H_0 false" $\frac{4020 - 4014}{60 / \sqrt{50}}$ $= 0.7071$ [0.712 from 4013.958] $1 - \Phi(0.7071)$ $= 0.240$ [0.238 from 4013.958]	M1dep depM1 A1✓ A1 M1 A1 6	Standardise 4020 and 4014 $\sqrt{}$, allow 60^2 , cc With $\sqrt{50}$ or 50 Completely correct LHS, $\sqrt{}$ on their c z -value in range [0.707, 0.712] Normal tables, answer < 0.5 Answer in range [0.2375, 0.2405]
	(iii) Smaller Smaller cv, better test etc	B1 B1 2	"Smaller" stated, no invalidating reason Plausible reason
	(iv) Smaller Smaller cv, larger prob of Type I etc	B1 B1 2	"Smaller" stated, no invalidating reason Plausible reason
	(v) No, parent distribution known to be normal	B2 2	"No" stated, convincing reason SR: If B0, "No", reason that is not invalidating: B1

Mark Scheme 4734
June 2006

1	Add two Poisson distributions With mean 17 $P(27) = e^{-17} 17^{27} / 27!$ or $P(\leq 27) - P(\leq 26)$ 0.00634 or 0.0063, 0.0064 from tables	M1 A1 M1 A1	4	Use formula or table M1A1 0.0052 from N(17,17)
2	$H_0: p_1 = p_2 = p_3 = p_4$, (H_1 : They are not all equal) Expected values under $H_0 = 150$ $\chi^2 = (12^2 + 23^2 + 15^2 + 20^2) / 150$ = 8.653 Critical value with 3 d.f. = 7.815 ($\chi^2 > 7.815$ so) reject H_0 and accept that proportions are different.	B1 B1 M1 B1 B1√	A1 6	Indication of equality of proportions At least one correct term Accept art 8.65 or 8.66 ft critical value
3	Assume population of differences has a normal distribution. or sample random $H_0: \mu_B - \mu_A = 0$, $H_1: \mu_B - \mu_A > 0$ $t = (23.43 - 22.84) / \sqrt{(0.548/10)}$ = 2.520 CV = 1.833 2.52 > CV so reject H_0 1.812, 1.734 Accept that there is evidence that mean time has reduced.	B1 B1 M1 B1 M1 A1√	A1 7	Either assumption. AEF Seen Allow from CV 2.262 (2-tail), ft wrong CV
4	(i) EITHER: $\int_{q_3}^4 \frac{1}{12} x dx = \frac{1}{4}$ or $\int_1^2 \frac{4}{3x^3} dx + \int_2^{q_3} \frac{1}{12} x dx = \frac{3}{4}$ [$x^2/24$] OR [$-2/(3x^2) + [x^2/24]$] ($16 - q_3^2$)/24 = 1/4 or $1/3 + q_3^2/24 = 3/4$ $q_3 = \sqrt{10}$ If they find F(x): M1A1, M1A1	M1* A1 dep *M1 A1	4	Either Form equation and attempt to solve Accept to 3 SF
(ii)	$E(X^2) = \int_1^2 \frac{4}{3x} dx + \int_2^4 \frac{x^3}{12} dx$ $E(X) = \int_1^2 \frac{4}{3x^2} dx + \int_2^4 \frac{x^2}{12} dx$ $\left[\frac{4}{3} \ln x \right]_1^2 + \left[\frac{x^4}{48} \right]_2^4$ $\left[\frac{-4}{3x} \right]_1^2 + \left[\frac{x^3}{36} \right]_2^4$ $a = E(X^2)/E(X)$ $a = 2.6659, 2.67$	M1 A1 A1 M1 A1	5	Either correct Or exact value, $(3 \ln 2)/5 + 9/4$ or equiv.

5	(i)	$(48 \times 72/150)$ or $(48/150)(72/150) \times 150$	M1 A1	2	Multiply and divide relevant values All correct
	(ii)	No, no expected value less than 5	B1	1	
	(iii)	H_0 : Volume and day are independent (H_1 : Volume and day are not independent) Critical value for 4 df = 13.28 Test statistic > 13.28, reject H_0 Accept that volume and day are not independent	B1 B1 M1 A1	4	Attributes specified
	(iv)	Choose Friday Highest volume	B1	B1 2	Not reference to E values
6	(i)	(a) No 0.43 belongs to relevant interval (b) Yes 0.43 is outside relevant interval	B1 B1 B1	3	Must be with reason
	(ii)	$H_0: p_R = p_T$, $H_1: p_R \neq p_T$ Estimate of $p = 74/165$ Variance estimate of difference $= \left(\frac{74}{165}\right)\left(\frac{91}{165}\right)\left(\frac{1}{80} + \frac{1}{85}\right)$ $z = (28/80 - 46/85)/\sigma_{\text{est}}$ $= -2.468$ Compare correctly with CV $-2.468 < -2.326$, or $2.468 > 2.326$ Reject H_0 and accept that the proportions differ on the island.	B1 B1 B1 M1 A1 A1 M1 A1	8	Proportions May be implied by later work Standardising Completely correct expression + or -, 2.47 Conclusion in context
7	(i)	$T_1 \sim N(2.2, 0.75^2)$, $T_2 \sim N(1.8, 0.70^2)$ Use $T_2 - \frac{1}{2} T_1$ normal $\mu = 0.7$ $\sigma^2 = 0.7^2 + \frac{1}{4} \times 0.75^2$ (0.630625) $(0 - \mu)/\sigma$ -0.881 Probability 0.189	M1 A1 A1 M1 A1	A1 6	Or $\frac{1}{2} T_1 - T_2$ From reasonable σ^2 not just sum + or -
	(ii)	Use sum of 5 Ts $\mu = 9.4$ $\sigma^2 = 2.5225$ $z = (10 - \mu)/\sigma$ Probability 0.6473, 0.647	M1 A1 A1 M1 A1	5	Standardising, must be σ
	(iii)	Calculation of variance	B1	1	

8	(i)	$s_B^2 = \frac{1}{49}(630.194 - \frac{176.35^2}{50})$	M1		Any equivalent formula
		$= 0.1675$	A1		May be implied by later work
		$H_0: \mu_B - \mu_A = 0, H_1: \mu_B - \mu_A > 0$		B1	aef
		$z = 0.115 / \sqrt{(0.049/40 + 0.1675/50)}$	M1		Standardising but not from pooled variance estimate
		$= 1.700$	A1		art 1.70
		$z > 1.645$, reject H_0 and accept that $\mu_B > \mu_A$	M1 A1	7	Compare correctly with 1.645 ft their calculated z
<hr/>					
--	(ii)	$z = 0.09 / \sqrt{(0.004575)}$	M1		Correct form
$= 1.331$		A1			
H_0 not rejected for $\alpha < 9.16$		M1 A1		Accept $< 9.2, \leq 9.2$. M1 for correct method for 9.2, A1 for inequality	
			4		
<hr/>					
<hr/>		(iii) (a) Not necessary		B1	Ignore any reason
<hr/>		(b) Not necessary since samples large enough for CLT to be applied (normality of sample means giving normality of difference)	M1		Mention of CLT implied by "sample large"
<hr/>			A1	3	Sample mean (approx) normal. (Do not award if population or sample said to be normal)

Mark Scheme 4735
June 2006

1	(i)	(a) True (b) False (c) True	B2	2	B0 for 0,1 correct, B1 for 2 correct, B2 for 3 correct.
	(ii)	Var(2X-Y)= 4Var(X)+Var(Y)-4Cov(X,Y) 6=11-4Cov(X,Y) Cov(X,Y)=5/4	M1 A1 A1	M1 4	Using formula Obtain cov cao
2		EITHER: sample is random OR twin pairs chosen independently H ₀ :m _F =m _S , H ₁ :m _F >m _S Use of B(60,0.5) Normal approx with μ=30, σ ² =15 EITHER: z=(36.5-30)/√15 =1.678 OR: CR is (X-30-0.5)/√15 > 1.645 X≥37 EITHER: 1.678> 1.645 OR: Sample value 37 in CR There is evidence that the first-born male twins are taller than the second -born twin in a majority of cases. OR: p-value: 0.0467 > 1.645 Completion NB: Exact Bin (60,0.5) p-value is 0.04623 from graphical calculator: full credit	B1 B1 M1 A2 A2 M1 A1 M1 A1	M1 A1 M1 M1 9	For both using medians Both Standardising A1 if correct apart from missing or wrong cc Setting up inequality A1 if correct apart from missing or wrong c.c. Correct comparison Conclusion in context
3	(i)	P(C)=P(C F)P(F)+P(C F')P(F') =0.98×0.05 + 0.04×0.95 0.087 AG	A1 A1	M1 3	Use of formula
	(ii)	P(F C)= =0.5632	M1A1 A1	3	art 0.563 or 49/87
	(iii)	P(F C')=P(C' F)P(F) / P(C') 0.02×0.05/0.913 [0.001095] 5000×above = 5.476., 5.48.	M1 M1A1	A1 4	Conditional prob. ft a conditional prob.

4	(i)	$M_X(t) = \int_a^b \frac{1}{b-a} e^{xt} dt$	M1	Correct integral with limits
		$= \left[\frac{e^{xt}}{(b-a)t} \right]_a^b$	B1	Correct integral
		$= \frac{e^{bt} - e^{at}}{(b-a)t}$ AG	A1	3






(ii)	Product of mgfs	M1	
	$\left(\frac{1-e^{-t}}{t} \right) \left(\frac{e^t-1}{t} \right)$	A1	2

(iii)	$M_S(t) = \left(\frac{e^{\frac{1}{2}t} - e^{-\frac{1}{2}t}}{t} \right)^2$	M1	Square of $M_Y(t)$
	$= (e^t - 2 + e^{-t})/t^2$	A1dep	Correctly shown
	mgfs of S and T are same	depA1	Correctly shown
	S and T have identical distributions	B1	4

5	(i)	${}^{13}C_4$ 715	M1 A1	2	Use of formula
		1234, 1235, 1236, 1237, 1245, 1246, 1345	B2 ft (i)		B1 for 5 or 6
		B1✓	3		7/715
	(iii)	Wilcoxon Rank Sum Test	B1		
		$H_0: m_X = m_Y, H_1: m_X \neq m_Y$	B1		Both, involving medians
		Use $P(R \leq 13)$	M1		
		$2 \times 7/715 \times 100 = 1.958\% < 2\%$	M1		Comparing correctly
		Reject H_0 , evidence of difference in medians at a significance level of (smaller than) 2%	A1	5	
		SR: If tables used,			
		B1B1 M1 for CV with correct comparison for rejection			
		M1 for rejection at 2% (not <)			
		Max 4/5			

6	(i)	$G'(t)=[0.8(1-0.2t)+0.16t]/(1-0.2t)^2$ $G'(t)=0.8/0.8^2=5/4$ AG	M1 A1 A1		Quotient or product rule	
				3		
						(ii)
		$G(t)=0.8t(1-0.2t)^{-1}$ $=0.8t(1+0.2t+0.04t^2+\dots)$ $P(Y=r)=0.8(0.2)^{r-1}$ $r=1,2,3,\dots$	M1 A1 A1 A1		Use binomial expansion At least 2 correct terms	
				4	OR from $G(0.8)$	
						(iii)
		EITHER: $Y \sim G(0.8)$ $\text{Var}(Y)=(1-0.8)/0.8^2$ $=0.3125$ OR: $G''(t)=0.32/(1-0.2t)^3$ Use $G''(1)+G'(1)-(G'(1))^2$ 0.3125	B1 M1 B1 M1 A1		Parameter not required A1	
				3		
						(iv)
		$G_T(t)=0.8^6 t^6 (1-0.2t)^{-6}$ $P(T \geq 8)=1-0.8^6(1+6 \times 0.2)$ $=0.42328$	B1 M1 A1		$(G_Y(t))^6$ Two terms in bracket art 0.423	
				3		
7	(i)	$E(X)=\frac{1}{2}(n+1)$ $\text{Var}(X)=\frac{1}{n}\sum r^2 - \frac{1}{4}(n+1)^2$ $=\frac{1}{6}(n+1)(2n+1) - \frac{1}{4}(n+1)^2$ $=\frac{1}{12}(n^2-1)$ AG	B1 M1 A1		Use of variance formula	
				4	Correctly obtained	
	(ii)	$E(N_1)=E(X_1)+E(X_2)-1$ $=\frac{1}{2}(n+1)+\frac{1}{2}(n+1)-1$ $=n$, (so N_1 is an unbiased estimator of n)	M1 A1			
				2		
	(iii)	$P(M=r)=$ EITHER: $P(X_1 < r, X_2 = r) + P(X_1 = r, X_2 < r)$ $=((r-1)/n)(1/(n-1)) + (1/n)(r-1)/(n-1)$ $=2(r-1)/[n(n-1)]$ AG, $r=2,3,4,\dots$ OR: Choose 1 from $r-1$ and 1 from 1 $\frac{r-1}{n} \times \frac{1}{n} \times \frac{1}{n} \times \frac{1}{n} \times \frac{1}{n} \times \frac{1}{n}$ $=(r-1)/[1/2 n(n-1)] = \text{AG}$	M1 A1 A1 M1 A1 A1			
				3		
	(iv)	$E(M) = \frac{2}{n(n-1)} \sum_{r=2}^n r(r-1)$ $=\frac{2}{3}(n+1)$ $N_2 = \frac{3}{2}M-1$	M1 A1√		A1 3	ft $E(M)$
	(v)	$\text{Var}(N_1) < \text{Var}(N_2)$ or equivalent $\frac{1}{6}(n^2-n-2) < \frac{9}{4}\text{Var}(M)$ $\text{Var}(M) > \frac{2}{27}(n^2-n-2)$	M1 A1 A1√			Stated or implied ft N_2
				3		

Mark Scheme 4736
June 2006

1	(i)	<p>2 4 3 3 2 5 4</p> <p>Box 1 2 4 2</p> <p>Box 2 3 3</p> <p>Box 3 5</p> <p>Box 4 4</p>	<p>M1</p> <p>A1 [2]</p>	<p>For packing these seven weights into boxes with no more than 8 kg total in each box</p> <p>For this packing</p>
	(ii)	<p>5 4 4 3 3 2 2</p> <p>Box 1 5 3</p> <p>Box 2 4 4</p> <p>Box 3 3 2 2</p>	<p>B1</p> <p>M1</p> <p>A1 [3]</p>	<p>For putting the weights into decreasing order (may be implied from packing)</p> <p>For packing the seven weights into three boxes with no more than 8 kg total in each box</p> <p>For this packing</p>
	(iii)	<p>15×2^2 = 60 seconds</p>	<p>M1</p> <p>A1 [2]</p>	<p>For a correct calculation</p> <p>For 60 or 60 seconds or 1 minute</p>
2	(i)	<div style="display: flex; justify-content: space-around; align-items: flex-start;"> <div style="text-align: center;">  <p>graph A</p> </div> <div style="text-align: center;">  <p>graph B</p> </div> <div style="text-align: center;">  <p>graph C</p> </div> </div> <p>other solutions:</p> <div style="display: flex; justify-content: space-around; align-items: center;">  <p>or</p>  </div>	<p>M1</p> <p>A1 [2]</p> <p>-----</p> <p>M1</p> <p>A1 [2]</p> <p>-----</p> <p>M1</p> <p>A1 [2]</p>	<p>Graphs may be in any order</p> <p>For a reasonable attempt</p> <p>For a graph that is topologically equivalent to one of these graphs</p> <p>For a different reasonable attempt</p> <p>For a graph that is topologically equivalent to one of these graphs</p> <p>For another different reasonable attempt</p> <p>For a graph that is topologically equivalent to one of these graphs</p>
	(ii)	<p>The graphs each have four odd nodes, but Eulerian graphs have no odd nodes.</p>	<p>B1 [1]</p>	<p>For any recognition that the nodes are not all even</p>

3	(i)	Travelling salesperson	B1 [1]	Identifying TSP by name
	(ii)	A – B – E – G – F – D – C – A	M1	For starting with A – B – E – G - ...
		130 (minutes)	A1	For this closed tour
		Shortest possible time ≤ 130 minutes	B1	For 130
			B1 [4]	For less than or equal to their time, with units
	(iii)	Order of connecting: B, E, G, F, D, C	B1	For a valid vertex order (or arc order) for their starting point
		<p>Lower bound = $10 + 15 + 95$ = 120 minutes</p>	M1	For a diagram or listing showing a tree connecting the vertices B, C, D, E, F and G but not A
			A1	For a diagram showing one of these trees (vertices must be labelled but arc weights are not needed)
			M1	For stating or using the total weight of their tree
			M1	For stating or using AB and AD or $10 + 15$
			A1 [6]	For 120 or calculating $25 +$ their 95, with units
	(iv)	A – B – E – G – F – C – D – A	M1	For a reasonable attempt
		or this in reverse	A1 [2]	For a valid tour of weight 125
				13

4	(i)	$x \leq 2$ $y \geq 1$ $y \leq 2x$ $x + y \leq 4$	B1 B1 B1 [4]	Strict inequalities used, penalise first time only All inequalities reversed, penalise first time only															
	(ii)	$(2, 1), (2, 2)$ $(\frac{1}{2}, 1)$ $(1\frac{1}{3}, 2\frac{2}{3})$	B1 B1 B1 [3]	Both of these This vertex in any exact form This vertex in any exact form or correct to 3 sf															
	(iii)	<table border="1"> <thead> <tr> <th>x</th> <th>y</th> <th>$P = x + 2y$</th> </tr> </thead> <tbody> <tr> <td>2</td> <td>1</td> <td>4</td> </tr> <tr> <td>2</td> <td>2</td> <td>6</td> </tr> <tr> <td>$\frac{1}{2}$</td> <td>1</td> <td>$2\frac{1}{2}$</td> </tr> <tr> <td>$1\frac{1}{3}$</td> <td>$2\frac{2}{3}$</td> <td>$6\frac{2}{3}$</td> </tr> </tbody> </table> $x = 1\frac{1}{3}, y = 2\frac{2}{3}$ (may be given in coordinate form) $P = 6\frac{2}{3}$	x	y	$P = x + 2y$	2	1	4	2	2	6	$\frac{1}{2}$	1	$2\frac{1}{2}$	$1\frac{1}{3}$	$2\frac{2}{3}$	$6\frac{2}{3}$	M1 A1 A1 [3]	Evidence of checking value at any vertex or using a sliding profit line Their x and y values at maximum in any exact form or correct to 3 sf Their maximum P value in any exact form or correct to 3 sf
	x	y	$P = x + 2y$																
	2	1	4																
2	2	6																	
$\frac{1}{2}$	1	$2\frac{1}{2}$																	
$1\frac{1}{3}$	$2\frac{2}{3}$	$6\frac{2}{3}$																	
(iv)	<table border="1"> <thead> <tr> <th>x</th> <th>y</th> <th>$Q = 2x - y$</th> </tr> </thead> <tbody> <tr> <td>2</td> <td>1</td> <td>3</td> </tr> <tr> <td>2</td> <td>2</td> <td>2</td> </tr> <tr> <td>$\frac{1}{2}$</td> <td>1</td> <td>0</td> </tr> <tr> <td>$1\frac{1}{3}$</td> <td>$2\frac{2}{3}$</td> <td>0</td> </tr> </tbody> </table> $Q = 0$ (x, y) can be any point on the line segment joining $(\frac{1}{2}, 1)$ and $(1\frac{1}{3}, 2\frac{2}{3})$	x	y	$Q = 2x - y$	2	1	3	2	2	2	$\frac{1}{2}$	1	0	$1\frac{1}{3}$	$2\frac{2}{3}$	0	M1 A1 A1 [3]	Evidence of checking value at any vertex or using a sliding profit line 0 (cao) The edge of the feasible region where $y = 2x$ No follow through	
x	y	$Q = 2x - y$																	
2	1	3																	
2	2	2																	
$\frac{1}{2}$	1	0																	
$1\frac{1}{3}$	$2\frac{2}{3}$	0																	
(v)	$P = Q \Rightarrow 2x - y = x + 2y$ $\Rightarrow x = 3y$ $y = \frac{1}{3}x$ lies entirely in the shaded region	M1 A1 A1 [3]	For considering $P = Q$, or equivalent For this line, or any equivalent reasoning For explanation of why there are no solutions																

16

5	(i)	$2x - 5y + 2z + s = 10$ $2x + 3z + t = 30$	B1 [1]	Slack variables used correctly																												
	(ii)	<table border="1"> <tr> <th>P</th> <th>x</th> <th>y</th> <th>z</th> <th>s</th> <th>t</th> <th></th> </tr> <tr> <td>1</td> <td>-1</td> <td>2</td> <td>3</td> <td>0</td> <td>0</td> <td>0</td> </tr> <tr> <td>0</td> <td>2</td> <td>-5</td> <td>2</td> <td>1</td> <td>0</td> <td>10</td> </tr> <tr> <td>0</td> <td>2</td> <td>0</td> <td>3</td> <td>0</td> <td>1</td> <td>30</td> </tr> </table>	P	x	y	z	s	t		1	-1	2	3	0	0	0	0	2	-5	2	1	0	10	0	2	0	3	0	1	30	M1 A1 [2]	<p>For overall structure correct, including two slack variable columns and column for RHS (condone omission of P column or labels)</p> <p>For a completely correct initial tableau, with no extra constraints added (condone variations in order of rows or columns)</p>
P	x	y	z	s	t																											
1	-1	2	3	0	0	0																										
0	2	-5	2	1	0	10																										
0	2	0	3	0	1	30																										
	(iii)	Pivot on x column since it is the only column with a negative value in the objective row $10 \div 2 = 5$ $5 < 15$ so pivot on this row $30 \div 2 = 15$	B1 B1 [2]	<p>For negative in objective row, top row, pay-off row, or equivalent</p> <p>For these two divisions shown</p>																												
	(iv)	New row 2 = row 2 \div 2 New row 1 = row 1 + new row 2 New row 3 = row 3 - 2 \times new row 2 <table border="1"> <tr> <td>1</td> <td>0</td> <td>-0.5</td> <td>4</td> <td>0.5</td> <td>0</td> <td>5</td> </tr> <tr> <td>0</td> <td>1</td> <td>-2.5</td> <td>1</td> <td>0.5</td> <td>0</td> <td>5</td> </tr> <tr> <td>0</td> <td>0</td> <td>5</td> <td>1</td> <td>-1</td> <td>1</td> <td>20</td> </tr> </table>	1	0	-0.5	4	0.5	0	5	0	1	-2.5	1	0.5	0	5	0	0	5	1	-1	1	20	B1 B1 [2] M1 M1 A1 [3]	<p>For dealing with the pivot row correctly</p> <p>For dealing with the other rows correctly</p> <p>May be coded by rows of table</p> <p>For updating their pivot row correctly</p> <p>For a reasonable attempt at updating other rows</p> <p>For correct values in tableau (condone consistent order of rows or columns). Do not follow through errors in initial tableau or pivot choice.</p>							
1	0	-0.5	4	0.5	0	5																										
0	1	-2.5	1	0.5	0	5																										
0	0	5	1	-1	1	20																										
		$x = 5, y = 0, z = 0$ $P = 5$ Not the maximum feasible value of P since there is still a negative value in the objective row	B1 B1 B1 [3]	<p>For reading off x, y and z from their tableau</p> <p>For reading off P from their tableau</p> <p>'No' seen or implied and a correct reason</p>																												

13


13

Mark Scheme 4737
June 2006

1	(i)	$4+4+8+7+6 = 29$ litres per second	B1 [1]	For 29
	(ii)	$4-1-2+3+3+5 = 12$ litres per second $0 - 5 - 4 + 3 + 0 + 5 = -1$ So minimum flow across cut is 0	M1 A1 M1 A1 [4]	For using upper and lower capacities correctly For showing how 12 (given) was worked out For a substantially correct calculation For 0, from an appropriate calculation
	(iii)	Flow in arc $CE \geq 2$ and flow in arc $CF \geq 3$, so at least 5 litres per second must flow into C At most 4 litres per second flow into A, of which at least 1 flows out to B and 2 flow out to E, so at most 1 litre per second can flow along AD	M1 A1 M1 A1 [4]	For any reasonable attempt (eg $CE = 2$, $CF = 3$) For correct reasoning For identifying ≤ 4 in and ≥ 3 out or equivalent For a correct conclusion
	(iv)	Either a diagram or a description of a flow of 11 litres per second. Arcs AD, AE, BE, CE, CF must all be at their minimum capacities.	M1 A1 A1 [3]	For a flow of 11 litres per second from S to T Flow satisfies all lower capacities Flow satisfies all upper capacities
	(v)	$11 \leq \text{maximum flow} \leq 12$	B1 B1 [2]	11 as lower bound 12 as upper bound (max flow = 12 \Rightarrow B0, B1)
				14

3	(i) 3 Y	M1 A1 [2]	For 3 (allow -3) For Y (cao)																				
(ii)	<p>$5 > 3, -2 > -4, 5 > -1$ and $6 > 0$ or using signs of differences +2, +2, +6, +6</p> <p>$3 > -2, -5 > -6, 1 > 0, 4 > 2$ or equivalent, or using differences</p> <p>Reduced matrix:</p> <table border="1" data-bbox="422 604 646 806"> <tr> <td></td><td colspan="3">Colin's strategy</td></tr> <tr> <td></td><td>W</td><td>X</td><td>Y</td></tr> <tr> <td>A</td><td>-1</td><td>4</td><td>-3</td></tr> <tr> <td>B</td><td>5</td><td>-2</td><td>5</td></tr> <tr> <td>D</td><td>-5</td><td>6</td><td>-4</td></tr> </table> <p>Rose's strategy</p>		Colin's strategy				W	X	Y	A	-1	4	-3	B	5	-2	5	D	-5	6	-4	<p>M1 A1</p> <p>M1 A1</p> <p>B1 [5]</p>	<p>For an appropriate comparison, or implied For all four comparisons seen</p> <p>For an appropriate comparison, or implied For all four comparisons seen</p> <p>For correct reduced matrix, with rows and columns labelled A, B, D and W, X, Y Cao</p>
	Colin's strategy																						
	W	X	Y																				
A	-1	4	-3																				
B	5	-2	5																				
D	-5	6	-4																				
(iii)	<p>Row minima are -3, -2, -5 Play-safe for Rose is B</p> <p>Column maxima are 5, 6, 5 Play-safes for Colin are W and Y</p> <p>Not stable</p>	<p>M1</p> <p>M1</p> <p>A1 [3]</p>	<p>Follow through their 3x3 reduced matrix For identifying row B</p> <p>For identifying columns W and Y</p> <p>For 'no' or 'not stable'</p>																				
(iv)	<p>5 is added throughout the matrix to make the entries non-negative. In this augmented reduced matrix, $9p_1 + 3p_2 + 11p_3$ is the expected number of points won by Rose when Colin plays strategy X</p>	<p>M1</p> <p>A1 [2]</p>	<p>For 'add 5' or equivalent</p> <p>For identifying that this is when Colin plays strategy X</p>																				
(v)	<p>$p_1 = \frac{7}{48}, p_2 = \frac{27}{48}, p_3 = \frac{14}{48}$ $\Rightarrow m \leq \frac{298}{48}$ (or $6\frac{5}{24}, 6.2083, 6.21$) in all three cases $\Rightarrow M = \frac{58}{48}$ (or $\frac{29}{24}, 1\frac{5}{24}, 1.2083, 1.21$)</p>	<p>M1</p> <p>A1 [2]</p>	<p>For attempting to evaluate m cao (in any appropriate form)</p>																				

4	(i)	<table><thead><tr><th>Activity</th><th>Duration</th><th>Immediate predecessors</th></tr></thead><tbody><tr><td>A</td><td>6</td><td>-</td></tr><tr><td>B</td><td>4</td><td>-</td></tr><tr><td>C</td><td>5</td><td>A</td></tr><tr><td>D</td><td>1</td><td>A, B</td></tr><tr><td>E</td><td>5</td><td>A, D</td></tr><tr><td>F</td><td>4</td><td>D</td></tr><tr><td>G</td><td>2</td><td>C, E, F</td></tr></tbody></table>	Activity	Duration	Immediate predecessors	A	6	-	B	4	-	C	5	A	D	1	A, B	E	5	A, D	F	4	D	G	2	C, E, F	B1	ANSWERED ON INSERT	
Activity	Duration	Immediate predecessors																											
A	6	-																											
B	4	-																											
C	5	A																											
D	1	A, B																											
E	5	A, D																											
F	4	D																											
G	2	C, E, F																											
			B1	For predecessors for activities A, B and C correct																									
			B1 [3]	For predecessors for activities D, F and G correct																									
				For predecessors for activity E correct																									
(ii)	<p>Minimum completion time = 14 hours Critical activities: A, D, E, G</p>	M1	For carrying out forward pass (no more than one independent error)																										
		A1	For all early event times correct																										
		M1	For carrying out backwards pass (no more than one independent error)																										
		A1	For all late event times correct																										
		B1 B1 [6]	For 14 For A, D, E, G only																										
(iii)	Increased by 2 (hours) Becomes 16 (hours)		B1	For stating that time increases by 2, or equivalent																									
(iv)	<p>Number of workers required = 3</p>	B1	For a resource histogram with no overhanging cells																										
		M1	For a reasonable attempt, fit their start times if possible																										
		A1	For a completely correct histogram (cao)																										
		B1 [4]	For 3 or follow through their histogram if possible																										
			14																										

5	(i)		M1 A1 [2]	ANSWERED ON INSERT For a substantially correct attempt For a completely correct bipartite graph																																																																								
	(ii)	$C-N \quad E-M \quad F-K$ $A-J \quad B-L \quad D-O$	M1 A1 [2]	For pairing $F-K, C-N, E-M$ For all correct (Diagram only \oplus M1, A0)																																																																								
	(iii)	<table border="1" data-bbox="220 815 740 1050"><tr><th></th><th>J</th><th>K</th><th>L</th><th>M</th><th>N</th><th>O</th></tr><tr><th>A</th><td>2</td><td>5</td><td>2</td><td>2</td><td>5</td><td>2</td></tr><tr><th>B</th><td>2</td><td>5</td><td>2</td><td>0</td><td>5</td><td>5</td></tr><tr><th>C</th><td>5</td><td>0</td><td>5</td><td>5</td><td>2</td><td>2</td></tr><tr><th>D</th><td>2</td><td>5</td><td>0</td><td>5</td><td>5</td><td>2</td></tr><tr><th>E</th><td>5</td><td>2</td><td>5</td><td>2</td><td>0</td><td>5</td></tr><tr><th>F</th><td>2</td><td>2</td><td>5</td><td>5</td><td>2</td><td>2</td></tr></table>		J	K	L	M	N	O	A	2	5	2	2	5	2	B	2	5	2	0	5	5	C	5	0	5	5	2	2	D	2	5	0	5	5	2	E	5	2	5	2	0	5	F	2	2	5	5	2	2	B1 B1 B1 [3]	For '5' in all the entries that should be 5 For '2' in all the entries that should be 2 For '0' in all the entries that should be 0																							
	J	K	L	M	N	O																																																																						
A	2	5	2	2	5	2																																																																						
B	2	5	2	0	5	5																																																																						
C	5	0	5	5	2	2																																																																						
D	2	5	0	5	5	2																																																																						
E	5	2	5	2	0	5																																																																						
F	2	2	5	5	2	2																																																																						
	(iv)	<p>Reduce rows</p> <table border="1" data-bbox="293 1106 740 1308"><tr><td>0</td><td>3</td><td>0</td><td>0</td><td>3</td><td>0</td></tr><tr><td>2</td><td>5</td><td>2</td><td>0</td><td>5</td><td>5</td></tr><tr><td>5</td><td>0</td><td>5</td><td>5</td><td>2</td><td>2</td></tr><tr><td>2</td><td>5</td><td>0</td><td>5</td><td>5</td><td>2</td></tr><tr><td>5</td><td>2</td><td>5</td><td>2</td><td>0</td><td>5</td></tr><tr><td>0</td><td>0</td><td>3</td><td>3</td><td>0</td><td>0</td></tr></table> <p>Columns are already reduced</p> <p>Or, reduce columns</p> <table border="1" data-bbox="293 1397 740 1597"><tr><td>0</td><td>5</td><td>2</td><td>2</td><td>5</td><td>0</td></tr><tr><td>0</td><td>5</td><td>2</td><td>0</td><td>5</td><td>3</td></tr><tr><td>3</td><td>0</td><td>5</td><td>5</td><td>2</td><td>0</td></tr><tr><td>0</td><td>5</td><td>0</td><td>5</td><td>5</td><td>0</td></tr><tr><td>3</td><td>2</td><td>5</td><td>2</td><td>0</td><td>3</td></tr><tr><td>0</td><td>2</td><td>5</td><td>5</td><td>2</td><td>0</td></tr></table> <p>Rows are already reduced</p> <p>Cannot cross out 0's using fewer than 6 lines so matching is complete</p> <p>$A-J \quad B-M \quad C-K \quad D-L \quad E-N$ $F-O$ $A-O \quad B-M \quad C-K \quad D-L \quad E-N$ $F-J$</p> <p>First matching: Fred and Jenny Second matching: Jenny and Olivia</p>	0	3	0	0	3	0	2	5	2	0	5	5	5	0	5	5	2	2	2	5	0	5	5	2	5	2	5	2	0	5	0	0	3	3	0	0	0	5	2	2	5	0	0	5	2	0	5	3	3	0	5	5	2	0	0	5	0	5	5	0	3	2	5	2	0	3	0	2	5	5	2	0	M1 A1 M1 A1 B1 B1 B1 B1 [8]	For a substantially correct attempt from their matrix For a correct reduction of rows and columns (or columns and rows) for their matrix For achieving a reduced cost matrix with a complete matching of zero cost (without unnecessary augmenting) 0's in correct cells (not ft) For this matching or ft their reduced cost matrix For this matching or ft their reduced cost matrix For the names for their first matching For the names for their second matching
0	3	0	0	3	0																																																																							
2	5	2	0	5	5																																																																							
5	0	5	5	2	2																																																																							
2	5	0	5	5	2																																																																							
5	2	5	2	0	5																																																																							
0	0	3	3	0	0																																																																							
0	5	2	2	5	0																																																																							
0	5	2	0	5	3																																																																							
3	0	5	5	2	0																																																																							
0	5	0	5	5	0																																																																							
3	2	5	2	0	3																																																																							
0	2	5	5	2	0																																																																							

Advanced GCE Mathematics (3890, 3892, 7890)
June 2006 Assessment Series

Unit Threshold Marks

Unit		Maximum Mark	a	b	c	d	e	u
4721	Raw	72	56	48	40	33	26	0
	UMS	100	80	70	60	50	40	0
4722	Raw	72	53	45	37	29	22	0
	UMS	100	80	70	60	50	40	0
4723	Raw	72	57	49	42	35	28	0
	UMS	100	80	70	60	50	40	0
4724	Raw	72	60	52	44	37	30	0
	UMS	100	80	70	60	50	40	0
4725	Raw	72	60	52	44	37	30	0
	UMS	100	80	70	60	50	40	0
4726	Raw	72	54	47	40	33	27	0
	UMS	100	80	70	60	50	40	0
4727	Raw	72	50	43	37	31	25	0
	UMS	100	80	70	60	50	40	0
4728	Raw	72	58	50	42	35	28	0
	UMS	100	80	70	60	50	40	0
4729	Raw	72	59	51	43	36	29	0
	UMS	100	80	70	60	50	40	0
4730	Raw	72	58	50	43	36	29	0
	UMS	100	80	70	60	50	40	0
4731	Raw	72	51	44	37	30	23	0
	UMS	100	80	70	60	50	40	0
4732	Raw	72	56	49	42	35	29	0
	UMS	100	80	70	60	50	40	0
4733	Raw	72	52	44	36	29	22	0
	UMS	100	80	70	60	50	40	0
4734	Raw	72	57	49	42	35	28	0
	UMS	100	80	70	60	50	40	0
4735	Raw	72	54	47	40	33	27	0
	UMS	100	80	70	60	50	40	0
4736	Raw	72	61	53	46	39	32	0
	UMS	100	80	70	60	50	40	0

4737	Raw	72	61	53	45	38	31	0
	UMS	100	80	70	60	50	40	0

Specification Aggregation Results

Overall threshold marks in UMS (i.e. after conversion of raw marks to uniform marks)

	Maximum Mark	A	B	C	D	E	U
3890	300	240	210	180	150	120	0
3891	300	240	210	180	150	120	0
3892	300	240	210	180	150	120	0
7890	600	480	420	360	300	240	0
7891	600	480	420	360	300	240	0
7892	600	480	420	360	300	240	0

The cumulative percentage of candidates awarded each grade was as follows:

	A	B	C	D	E	U	Total Number of Candidates
3890	31.0	46.3	61.2	73.5	84.2	100	12438
3891	0	0	0	100	100	100	1
3892	60.6	76.8	89.2	95.3	97.6	100	1109
7890	46.9	67.7	81.9	91.5	97.6	100	9525
7891	50.0	75.0	87.5	87.5	100	100	8
7892	59.9	80.2	89.4	95.5	98.6	100	1428

For a description of how UMS marks are calculated see;
www.ocr.org.uk/OCR/WebSite/docroot/understand/ums.jsp

Statistics are correct at the time of publication